



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Elzovantinib

Cat. No.:	HY-111787
CAS No.:	2271119-26-5
Molecular Formula:	C ₂₀ H ₂₀ FN ₇ O ₂
Molecular Weight:	409.42
Target:	Src; c-Met/HGFR; c-Fms
Pathway:	Protein Tyrosine Kinase/RTK
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 25 mg/mL (61.06 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		2.4425 mL	12.2124 mL	24.4248 mL
		5 mM		0.4885 mL	2.4425 mL	4.8850 mL
		10 mM		0.2442 mL	1.2212 mL	2.4425 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (5.08 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (5.08 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	Elzovantinib (TPX-0022) is an oral-active inhibitor of SRC, MET and c-FMS, with IC ₅₀ values of 0.12 nM, 0.14 nM and 0.76 nM for SRC, MET and c-FMS respectively ^[1] .
IC ₅₀ & Target	IC ₅₀ : 0.12 nM (SRC), 0.14 nM (MET), 0.76 nM (c-FMS) ^[1] .
In Vitro	Elzovantinib (TPX-0022) causes the suppression of MET autophosphorylation as well as the downstream STAT3, ERK and AKT phosphorylation at IC ₅₀ values of around 1-3 nM in SNU-5 and MKN-45 cell lines ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	Elzovantinib (TPX-0022; p.o., BID, 13 days) treatment results in an 85% tumor regression and no body weight loss is observed

after 21 days treatment in mice^[1].

Elzovantinib (p.o., BID, 10 days) demonstrates the ability to inhibit tumor growth at 44% and 67% at the dose of 5 mg/kg, BID and 15 mg/kg, BID, respectively in SCID/Beige mice^[1].

Elzovantinib inhibits MET activity in MKN-45 tumors following oral administration in mice^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Mice bearing LU2503 tumors patient derived xenograft (PDX) NSCLC model ^[1] .
Dosage:	15 mg/kg.
Administration:	PO, BID (twice daily) for 13 days.
Result:	Resulted in an 85% tumor regression and no body weight loss was observed after 21 days treatment.

Animal Model:	SCID/Beige mice bearing Ba/F3 ETV6-CSF1R tumors with average tumor size of ~180 mm ³ ^[1] .
Dosage:	5 and 15 mg/kg.
Administration:	PO, BID (twice daily) for 10 days.
Result:	Demonstrated the ability to inhibit tumor growth at 44% and 67% at the dose of 5 mg/kg, BID and 15 mg/kg, BID, respectively.

REFERENCES

[1]. WO 2019023417 A1.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA