

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



## **Product** Data Sheet

### eCF506-d5

Molecular Weight: 515.66

Target: Src; Isotope-Labeled Compounds

Pathway: Protein Tyrosine Kinase/RTK; Others

Storage: Please store the product under the recommended conditions in the Certificate of

Analysis.

#### **BIOLOGICAL ACTIVITY**

Description	eCF506- $d_5$ is deuterated labeled eCF506 (HY-112096). eCF506 is a highly potent and orally bioavailable inhibitor of the non-receptor tyrosine kinase Src with an IC <sub>50</sub> of less than 0.5 nM.
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . eCF506 induces a very potent antiproliferative effect in both MCF7 and MDA-MB-231 cells. eCF506 inhibits phosphorylation of SRC and FAK at low nanomolar levels, with complete inhibition observed at 100 nM. eCF506 significantly reduces cell motility at 10 nM as early as 6 h into the study, with equivalent efficacy to dasatinib. eCF506 exclusively inhibits SFK, with subnanomolar IC $_{50}$ values against SRC and YES (IC $_{50}$ =0.5, 2.1 nM). It is important to highlight that eCF506 displays a vast difference in activity (>950-fold difference) between ABL and its primary target SRC <sup>[2]</sup> .
In Vivo	eCF506 shows a moderate oral bioavailability (25.3%). A significant reduction of phospho-SRC <sup>Y416</sup> is observed in the xenograft sections from mice treated with eCF506 relative to the untreated animal controls <sup>[2]</sup> .  MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### **REFERENCES**

[1]. Fraser C, et al. Rapid Discovery and Structure-Activity Relationships of Pyrazolopyrimidines That Potently Suppress Breast Cancer Cell Growth via SRC Kinase Inhibition with Exceptional Selectivity over ABL Kinase. J Med Chem. 2016 May 26;59(10):4697-710.

[2]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA