



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

6-(γ,γ -Dimethylallylamino)purine-d₆

Cat. No.:	HY-112103S
CAS No.:	175733-28-5
Molecular Formula:	C ₁₀ H ₇ D ₆ N ₅
Molecular Weight:	209.28
Target:	Isotope-Labeled Compounds
Pathway:	Others
Storage:	Powder -20°C 3 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 60 mg/mL (286.70 mM; Need ultrasonic and warming)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		4.7783 mL	23.8914 mL	47.7829 mL
	5 mM		0.9557 mL	4.7783 mL	9.5566 mL
	10 mM		0.4778 mL	2.3891 mL	4.7783 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

6-(γ,γ -Dimethylallylamino)purine-d₆ is the deuterium labeled [6-\(\$\gamma,\gamma\$ -Dimethylallylamino\)purine](#) (HY-112103). 6-(γ,γ -Dimethylallylamino)purine is a plant growth substance[1][2].

In Vitro

Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs^[1].

Derivative UV spectroscopic data show that the plant growth substances 6-(γ,γ -Dimethylallylamino)purine (N6-(delta 2-isopentenyl) adenine, i6Ade) and indolylacetic acid (IAA) can bind to the yeast alcohol dehydrogenase (ADH) and affect coenzyme-enzyme binding. At fixed ethanol concentrations (27.8 and 111.1 mM) and varying NAD⁺ concentrations (0.033-2 mM), as well as at fixed levels of coenzyme (0.67 and 2 mM), and at varying concentrations of ethanol (1.4-111.1 mM), the rate of ethanol oxidation is significantly inhibited by i6Ade and IAA. The kinetics of the ADH reaction is affected by two inhibition constants (K_i and K_i') which correspond to the dissociation constants of complexes EI and ESI, respectively. For i6Ade the K_i=0.52±0.06 mM and K_i'=0.74±0.07 mM, and for IAA the K_i=0.88±0.03 mM and K_i'=0.99±0.02 mM^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Zikmanis P, et al. Indolylacetic acid and N6-(delta 2-isopentenyl) adenine affect NADH binding to yeast alcohol dehydrogenase and inhibit in vitro the enzymatic oxidation of ethanol. Biofactors. 1990 Oct;2(4):237-40.
- [2]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019;53(2):211-223.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA