



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



NF449 octasodium

Cat. No.:	HY-112461A
CAS No.:	627034-85-9
Molecular Formula:	C ₄₁ H ₂₄ N ₆ Na ₈ O ₂₉ S ₈
Molecular Weight:	1505.09
Target:	P2X Receptor
Pathway:	Membrane Transporter/Ion Channel
Storage:	-20°C, stored under nitrogen
	* In solvent : -80°C, 6 months; -20°C, 1 month (stored under nitrogen)



SOLVENT & SOLUBILITY

In Vitro

H₂O : 10 mg/mL (6.64 mM; Need ultrasonic and warming)
DMSO : 10 mg/mL (6.64 mM; ultrasonic and warming and heat to 60°C)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		0.6644 mL	3.3221 mL	6.6441 mL
	5 mM		0.1329 mL	0.6644 mL	1.3288 mL
	10 mM		---	---	---

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

NF449 octasodium is a highly potent P2X₁ receptor antagonist, with IC₅₀s of 0.28, 0.69, and 120 nM for rP2X₁, rP2X₁₊₅, P2X₂₊₃, respectively. NF449 octasodium is a G_{sα}-selective G Protein antagonist. NF449 octasodium suppresses the rate of GTP[γS] binding to G_{sα-s}, inhibits the stimulation of adenylyl cyclase activity, and blocks the coupling of β-adrenergic receptors to G_s [1][2].

IC₅₀ & Target

p2x1 Receptor

In Vitro

NF449 octasodium suppressed the rate of GTP[γS] binding to rG_{sα-s} while barely affecting binding to rG_{iα-1} (IC₅₀=140 nM), inhibits stimulation of adenylyl cyclase activity in S49 cyc membranes (deficient in endogenous Gsa) by exogenously added Gsa-s, and blocks the coupling of β-adrenergic receptors to G_s (EC₅₀=7.9 μM) [2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

At a dose of 10 mg/kg, NF449 octasodium inhibits the ex vivo aggregation triggered by 5 g/ml collagen in WT mouse platelets without affecting that induced by 5 μM ADP. At a higher dose (50 mg/kg), NF449 octasodium inhibits ex vivo platelet aggregation in response to not only 10 g/ml collagen but also 5 M ADP, indicating nonselective inhibition of the P2Y1 and/or P2Y12 receptor [3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Vet Parasitol. 2022 Nov 19;312:109841.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Rettinger J, et al. Profiling at recombinant homomeric and heteromeric rat P2X receptors identifies the suramin analogue NF449 as a highly potent P2X1 receptor antagonist. *Neuropharmacology*. 2005;48(3):461-468.
- [2]. Hechler B, et al. Inhibition of platelet functions and thrombosis through selective or nonselective inhibition of the platelet P2 receptors with increasing doses of NF449 [4,4',4'',4'''-(carbonylbis(imino-5,1,3-benzenetriylbis-(carbonylimino)))tetrakis-benzene-1,3-disulfonic acid octasodium salt]. *J Pharmacol Exp Ther*. 2005;314(1):232-243.
- [3]. Hohenegger M, et al. Gsalpha-selective G protein antagonists. *Proc Natl Acad Sci U S A*. 1998;95(1):346-351.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA