



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PF-06747711

Cat. No.:	HY-112706
CAS No.:	1892576-58-7
Molecular Formula:	C ₂₆ H ₂₆ F ₃ N ₅ O ₂
Molecular Weight:	497.51
Target:	ROR
Pathway:	Metabolic Enzyme/Protease; Vitamin D Related/Nuclear Receptor
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 41.67 mg/mL (83.76 mM; Need ultrasonic)				
	Preparing Stock Solutions	Solvent Concentration Mass	1 mg	5 mg	10 mg
		1 mM	2.0100 mL	10.0500 mL	20.1001 mL
		5 mM	0.4020 mL	2.0100 mL	4.0200 mL
		10 mM	0.2010 mL	1.0050 mL	2.0100 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (4.18 mM); Clear solution				
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (4.18 mM); Clear solution				
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (4.18 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	PF-06747711 is a potent, selective, and orally active retinoic acid receptor-related orphan C2 (RORC2, also known as RORγt) inverse agonist, with an IC ₅₀ of 4.1 nM. Anti-skin inflammatory activity ^[1] .
IC ₅₀ & Target	IC50: 4.1 nM (RORC2) ^[1]
In Vitro	PF-06747711 (Compound 66) reduces IL-17 production by human Th17 cells with an IC ₅₀ of 9.5 nM ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

PF-06747711 (10, 30, and 100 mg/kg, p.o., daily over 5 days) inhibits ear swelling in a dose-dependent manner in mice^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	8–10 week old female Balb/c mice ^[1]
Dosage:	10, 30, and 100 mg/kg
Administration:	P.O. daily over 5 days
Result:	Inhibited ear swelling, and caused a maximum inhibition of 46% at 100 mg/kg.

REFERENCES

[1]. Schnute ME, et al. Discovery of 3-Cyano- N-(3-(1-isobutylpiperidin-4-yl)-1-methyl-4-(trifluoromethyl)-1 H-pyrrolo[2,3- b]pyridin-5-yl)benzamide: A Potent, Selective, and Orally Bioavailable Retinoic Acid Receptor-Related Orphan Receptor C2 Inverse Agonist. J Med Chem. 2018 Sep 9.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA