

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



## HC-056456

Cat. No.: HY-112729 CAS No.: 7733-96-2 Molecular Formula:  $C_{12}H_{6}N_{2}O_{4}S_{2}$ Molecular Weight: 306.32

Target: Calcium Channel

Pathway: Membrane Transporter/Ion Channel; Neuronal Signaling

Storage: Powder -20°C 3 years

4°C 2 years -80°C 2 years

In solvent

-20°C 1 year

**Product** Data Sheet

#### **SOLVENT & SOLUBILITY**

In Vitro

DMSO: 125 mg/mL (408.07 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	3.2646 mL	16.3228 mL	32.6456 mL
	5 mM	0.6529 mL	3.2646 mL	6.5291 mL
	10 mM	0.3265 mL	1.6323 mL	3.2646 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (6.79 mM); Clear solution
- 2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (6.79 mM); Clear solution

### **BIOLOGICAL ACTIVITY**

Description	HC-056456 is an effective but not perfectly-selective blocker of CatSper channels. The $[Na^+]_i$ rise is slowed by HC-056456 (IC $_{50}^{-3} \mu M$ ).
IC <sub>50</sub> & Target	CatSper <sup>[1]</sup>
In Vitro	HC-056456 similarly slows the rise of $[Ca^{2+}]_i$ that is evoked by alkaline depolarization and reported by fura-2. HC-056456 also selectively and reversibly decreased CatSper currents recorded from patch-clamped sperm. HC-056456 produces a pharmacological phenocopy of the CatSper-null sperm. Acute application of HC-056456 causes rapid loss of flagellar waveform asymmetry from hyperactivated sperm, indicating that continued entry of Ca <sup>2+</sup> through CatSper channels is required to maintain hyperactivation. HC-056456 selectively and reversibly blocks CatSper currents. The specificity and

reversibility of the blockade of CatSper-dependent currents by HC-056456 is examined by using patch clamp recordings. The observed current is blocked slightly more than 50% by 20  $\mu$ M HC-056456 (estimated IC<sub>50</sub> near 15  $\mu$ M). In concept, it remains possible that CatSper channel heterogeneity explains residual HC-056456-resistant current. The action of HC-056456 on KSper channels, the other major cation channel observed in patch-clamped sperm, is also examined. Subsequent application of 50  $\mu$ M HC-056456 results in partial blockade of this current. For HC-056456 action on KSper an IC<sub>50</sub> near 40  $\mu$ M is estimated<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### **CUSTOMER VALIDATION**

- Comp Biochem Physiol A Mol Integr Physiol. 2020 Mar;241:110634.
- Comp Biochem Physiol A Mol Integr Physiol. 2020 Mar;241:110634.

See more customer validations on www.MedChemExpress.com

#### **REFERENCES**

[1]. Carlson AE, et al. Pharmacological targeting of native CatSper channels reveals a required role in maintenance of sperm hyperactivation. PLoS One. 2009 Aug 31;4(8):e6844.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

 $\hbox{E-mail: } tech@MedChemExpress.com$ 

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA