



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

C-176

Cat. No.:	HY-112906
CAS No.:	314054-00-7
Molecular Formula:	C ₁₁ H ₇ IN ₂ O ₄
Molecular Weight:	358.09
Target:	STING
Pathway:	Immunology/Inflammation
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : 62.5 mg/mL (174.54 mM; Need ultrasonic)
 Ethanol : 2.5 mg/mL (6.98 mM; ultrasonic and warming and heat to 60°C)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.7926 mL	13.9630 mL	27.9259 mL
	5 mM		0.5585 mL	2.7926 mL	5.5852 mL
	10 mM		0.2793 mL	1.3963 mL	2.7926 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: Cremophor EL
Solubility: 10 mg/mL (27.93 mM); Clear solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: 1.67 mg/mL (4.66 mM); Suspended solution; Need ultrasonic and warming
- Add each solvent one by one: 10% EtOH >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 0.25 mg/mL (0.70 mM); Clear solution
- Add each solvent one by one: 10% EtOH >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 0.25 mg/mL (0.70 mM); Clear solution
- Add each solvent one by one: 10% EtOH >> 90% corn oil
Solubility: 0.25 mg/mL (0.70 mM); Suspended solution; Need ultrasonic

BIOLOGICAL ACTIVITY

Description

C-176 is a selective and blood-brain barrier permeable STING inhibitor. C-176 covalently targets transmembrane cysteine residue 91 and thereby blocking activation-induced palmitoylation of STING^{[1][2]}.

IC ₅₀ & Target	STING ^[1] .								
In Vitro	C-176 strongly reduces STING-mediated, but not RIG-I- or TBK1-mediated, IFN β reporter activity. Pretreatment with C-176 markedly reduce the CMA-mediated induction of serum levels of type I IFNs and IL-6 ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.								
In Vivo	C-176 (750/375 nmol C-176 per mouse in 200 μL corn oil) significantly reduces the CMA-mediated induction of serum levels of type I IFNs and IL-6., without significant toxicity^[1]. C-176 results in a significant reduction in serum levels of type I IFNs and in a strong suppression of inflammatory parameters in the heart, with no evident signs of overt toxicity Trex1^{-/-} mice^[1]. C-176 demonstrates marked amelioration of various signs of systemic inflammation in Trex1^{-/-} mice^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only. <table> <tr> <td>Animal Model:</td><td>WT type mice.</td></tr> <tr> <td>Dosage:</td><td>750/375 nmol C-176 per mouse in 200 μL corn oil (~1.34/0.67 mg/mL).</td></tr> <tr> <td>Administration:</td><td>Intraperitoneally, once.</td></tr> <tr> <td>Result:</td><td>Significantly reduced Serum levels of type I IFNs and IL-6.</td></tr> </table>	Animal Model:	WT type mice.	Dosage:	750/375 nmol C-176 per mouse in 200 μ L corn oil (~1.34/0.67 mg/mL).	Administration:	Intraperitoneally, once.	Result:	Significantly reduced Serum levels of type I IFNs and IL-6.
Animal Model:	WT type mice.								
Dosage:	750/375 nmol C-176 per mouse in 200 μ L corn oil (~1.34/0.67 mg/mL).								
Administration:	Intraperitoneally, once.								
Result:	Significantly reduced Serum levels of type I IFNs and IL-6.								

CUSTOMER VALIDATION

- Bioact Mater. 2022 Dec 9;24:37-53.
- Nat Commun. 2023 May 23;14(1):2950.
- Nat Commun. 2023 May 26;14(1):3050.
- Neuron. 2022 Nov 4;S0896-6273(22)00961-8.
- J Clin Invest. 2021 Oct 15;131(20):e136329.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Zhang LM, et al. STING mediates neuroinflammatory response by activating NLRP3-related pyroptosis in severe traumatic brain injury. J Neurochem. 2022 Sep;162(5):444-462.
- [2]. Haag SM, et al. Targeting STING with covalent small-molecule inhibitors. Nature. 2018 Jul;559(7713):269-273.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA