



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GW284543

Cat. No.:	HY-114189
CAS No.:	790186-68-4
Molecular Formula:	C ₂₃ H ₂₀ N ₂ O ₃
Molecular Weight:	372.42
Target:	MEK
Pathway:	MAPK/ERK Pathway
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 125 mg/mL (335.64 mM; Need ultrasonic)				
	Preparing Stock Solutions	Solvent Concentration Mass	1 mg	5 mg	10 mg
		1 mM	2.6851 mL	13.4257 mL	26.8514 mL
		5 mM	0.5370 mL	2.6851 mL	5.3703 mL
		10 mM	0.2685 mL	1.3426 mL	2.6851 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (5.59 mM); Clear solution				
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (5.59 mM); Clear solution				
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (5.59 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	GW284543 (UNC10225170) is a selective MEK5 inhibitor. GW284543 reduces pERK5, and decreases endogenous MYC protein [1].
IC ₅₀ & Target	MEK5
In Vitro	GW284543 (10 and 20 μM; 6 h) treatment inhibits MEK5 in a dose-dependent manner and reduces pERK5 ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Western Blot Analysis^[1]

Cell Line:	MIA PaCa-2 cells
Concentration:	10 and 20 μ M
Incubation Time:	6 hours
Result:	Reduced phosphorylation of ERK5, and decreased endogenous MYC protein.

REFERENCES

[1]. Vaseva AV, et al. KRAS Suppression-Induced Degradation of MYC Is Antagonized by a MEK5-ERK5 Compensatory Mechanism. Cancer Cell. 2018 Nov 12;34(5):807-822.e7.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA