



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## NXT629

|                    |                                                                                  |
|--------------------|----------------------------------------------------------------------------------|
| Cat. No.:          | HY-114263                                                                        |
| CAS No.:           | 1454925-59-7                                                                     |
| Molecular Formula: | C <sub>35</sub> H <sub>39</sub> N <sub>5</sub> O <sub>3</sub> S                  |
| Molecular Weight:  | 609.78                                                                           |
| Target:            | PPAR                                                                             |
| Pathway:           | Cell Cycle/DNA Damage                                                            |
| Storage:           | Powder    -20°C    3 years<br>In solvent   -80°C    6 months<br>-20°C    1 month |



### SOLVENT & SOLUBILITY

|                                                                               |                                                                                                                                           |                                                                      |           |           |            |
|-------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------|-----------|-----------|------------|
| In Vitro                                                                      | DMSO : 125 mg/mL (204.99 mM; Need ultrasonic)                                                                                             |                                                                      |           |           |            |
|                                                                               | Preparing Stock Solutions                                                                                                                 | <div><div>Solvent</div><div>Concentration</div><div>Mass</div></div> | 1 mg      | 5 mg      | 10 mg      |
|                                                                               |                                                                                                                                           | 1 mM                                                                 | 1.6399 mL | 8.1997 mL | 16.3994 mL |
|                                                                               |                                                                                                                                           | 5 mM                                                                 | 0.3280 mL | 1.6399 mL | 3.2799 mL  |
|                                                                               |                                                                                                                                           | 10 mM                                                                | 0.1640 mL | 0.8200 mL | 1.6399 mL  |
| Please refer to the solubility information to select the appropriate solvent. |                                                                                                                                           |                                                                      |           |           |            |
| In Vivo                                                                       | 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline<br>Solubility: ≥ 2.08 mg/mL (3.41 mM); Clear solution |                                                                      |           |           |            |
|                                                                               | 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)<br>Solubility: ≥ 2.08 mg/mL (3.41 mM); Clear solution            |                                                                      |           |           |            |
|                                                                               | 3. Add each solvent one by one: 10% DMSO >> 90% corn oil                                                                                  |                                                                      |           |           |            |
|                                                                               | Solubility: ≥ 2.08 mg/mL (3.41 mM); Clear solution                                                                                        |                                                                      |           |           |            |

### BIOLOGICAL ACTIVITY

|                           |                                                                                                                                                                                                                                                                                                                                                                                                                                           |                                    |                                     |                                    |
|---------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------|-------------------------------------|------------------------------------|
| Description               | NXT629 is a potent, selective, and competitive PPAR-α antagonist, with an IC <sub>50</sub> of 77 nM for human PPARα, shows high selectivity over other nuclear hormone receptor, such as PPARδ, PPARγ, ERβ, GR and TRβ, IC <sub>50</sub> s are 6.0, 15, 15.2, 32.5 and >100 μM, respectively <sup>[1]</sup> . NXT629 has potent anti-tumor activity and inhibits experimental metastasis of cancer cell in animal models <sup>[2]</sup> . |                                    |                                     |                                    |
| IC <sub>50</sub> & Target | hPPARα<br>77 nM (IC <sub>50</sub> )                                                                                                                                                                                                                                                                                                                                                                                                       | hPPARδ<br>6 μM (IC <sub>50</sub> ) | hPPARγ<br>15 μM (IC <sub>50</sub> ) | ERβ<br>15.2 μM (IC <sub>50</sub> ) |
|                           | GR                                                                                                                                                                                                                                                                                                                                                                                                                                        |                                    |                                     |                                    |

|                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                 | 32.5 $\mu$ M (IC <sub>50</sub> )                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| <b>In Vitro</b> | <p>NXT629 (Compound 33) is a potent, selective PPAR-<math>\alpha</math> antagonist, with an IC<sub>50</sub> of 77 nM for human PPAR<math>\alpha</math>, shows high selectivity over other nuclear hormone receptor, such as PPAR<math>\delta</math>, PPAR<math>\gamma</math>, Er<math>\beta</math>, GR and TR<math>\beta</math>, IC<sub>50</sub>s are 6.0, 15, 15.2, 32.5 and &gt;100 <math>\mu</math>M, respectively<sup>[1]</sup>. NXT629 also competitively inhibits mouse PPAR<math>\alpha</math>, PPAR<math>\beta/\delta</math> and PPAR<math>\gamma</math>, with IC<sub>50</sub>s of 2.3, 35.1, 6.9 <math>\mu</math>M, respectively<sup>[2]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>          |
| <b>In Vivo</b>  | <p>NXT629 (Compound 33; 30 mg/kg, i.p.) exhibits good pharmacokinetics in mouse, and significantly decreases Fgf21 (Fibroblast growth factor 21), a PPAR<math>\alpha</math> target gene in fasted mice<sup>[1]</sup>.</p> <p>NXT629 has poor oral bioavailability in mice and rats. NXT629 (30 mg/kg, i.p., daily for 6 weeks) delays growth of subcutaneous SKOV-3 tumors in nude mice, inhibits growth of subcutaneous B16F10 tumors in C57Bl/6 mice. NXT629 (30 mg/kg, i.p.) is weakly anti-angiogenic against FGF-induced angiogenesis. NXT629 (3, 30 mg/kg, i.p.) inhibits experimental metastasis of B16F10 melanoma cells to the mouse lung<sup>[2]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> |

## CUSTOMER VALIDATION

- Nat Commun. 2021 Dec 2;12(1):7031.
- Cells. 2022 May 10;11(10):1597.
- Arch Physiol Biochem. 2022 Jan 21;1-18.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Bravo Y, et al. Identification of the first potent, selective and bioavailable PPAR $\alpha$  antagonist. Bioorg Med Chem Lett. 2014 May 15;24(10):2267-72.

[2]. Stebbins KJ, et al. In vitro and in vivo pharmacology of NXT629, a novel and selective PPAR $\alpha$  antagonist. Eur J Pharmacol. 2017 Aug 15;809:130-140.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA