



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## TCH-165

|                    |                                                               |       |         |
|--------------------|---------------------------------------------------------------|-------|---------|
| Cat. No.:          | HY-120722                                                     |       |         |
| CAS No.:           | 1446350-60-2                                                  |       |         |
| Molecular Formula: | C <sub>39</sub> H <sub>37</sub> N <sub>3</sub> O <sub>3</sub> |       |         |
| Molecular Weight:  | 595.73                                                        |       |         |
| Target:            | Proteasome                                                    |       |         |
| Pathway:           | Metabolic Enzyme/Protease                                     |       |         |
| Storage:           | Powder                                                        | -20°C | 3 years |
|                    |                                                               | 4°C   | 2 years |
|                    | In solvent                                                    | -80°C | 2 years |
|                    |                                                               | -20°C | 1 year  |



### SOLVENT & SOLUBILITY

|                                                                               |                                                                                                                                            |                                                       |      |           |           |            |
|-------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------|------|-----------|-----------|------------|
| In Vitro                                                                      | DMSO : 250 mg/mL (419.65 mM; Need ultrasonic)                                                                                              |                                                       |      |           |           |            |
|                                                                               | Preparing Stock Solutions                                                                                                                  | <div><div>Solvent</div><div>Concentration</div></div> | Mass | 1 mg      | 5 mg      | 10 mg      |
|                                                                               |                                                                                                                                            | 1 mM                                                  |      | 1.6786 mL | 8.3931 mL | 16.7861 mL |
|                                                                               |                                                                                                                                            | 5 mM                                                  |      | 0.3357 mL | 1.6786 mL | 3.3572 mL  |
|                                                                               |                                                                                                                                            | 10 mM                                                 |      | 0.1679 mL | 0.8393 mL | 1.6786 mL  |
| Please refer to the solubility information to select the appropriate solvent. |                                                                                                                                            |                                                       |      |           |           |            |
| In Vivo                                                                       | 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline<br>Solubility: ≥ 6.25 mg/mL (10.49 mM); Clear solution |                                                       |      |           |           |            |
|                                                                               | 2. Add each solvent one by one: 10% DMSO >> 90% corn oil<br>Solubility: ≥ 6.25 mg/mL (10.49 mM); Clear solution                            |                                                       |      |           |           |            |

### BIOLOGICAL ACTIVITY

|                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|---------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Description               | TCH-165 is a small molecule modulator of proteasome assembly, which increases 20S levels and facilitates 20S-mediated protein degradation <sup>[1]</sup> .                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| IC <sub>50</sub> & Target | Proteasome assembly <sup>[1]</sup>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| In Vitro                  | <p>TCH-165 (0.01-10 μM; 72 hours; RPMI8226 and U87MG cells) treatment inhibits cell growth of RPMI8226 and U87MG cells with IC<sub>50</sub> of 1.6 μM and 2.4 μM, respectively<sup>[1]</sup>.</p> <p>?TCH-165 (0-10 μM; 24 hours; HEK293T cells) treatment enhances ODC degradation is blocked by BTZ indicated that this event is proteasome-mediated. TCH-165 enhances proteolytic degradation in a concentration-dependent manner<sup>[1]</sup>.</p> <p>?TCH-165 enhances the chymotrypsin-like (CT-L), trypsin-like (Tryp-L) and caspase-like (Casp-L) activities with EC<sub>50</sub>s of 4.2 μ</p> |

M, 3.2  $\mu$ M and 4.7  $\mu$ M, respectively<sup>[1]</sup>.

?TCH-165 enhances 20S-mediated degradation of IDPs,  $\alpha$ -syn, and tau in vitro, and does not induce the degradation of structured proteins such as GAPDH<sup>[1]</sup>.

?TCH-165-treated cells display a decrease in the assembled 26S and an increase in the 20S proteasome. TCH-165 regulates the dynamic equilibrium between the 20S and 26S proteasome complexes, favoring 20S-mediated protein degradation<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Cell Viability Assay<sup>[1]</sup>

|                  |                                                                                                                       |
|------------------|-----------------------------------------------------------------------------------------------------------------------|
| Cell Line:       | RPMI8226 and U87MG cells                                                                                              |
| Concentration:   | 0.01-10 $\mu$ M                                                                                                       |
| Incubation Time: | 72 hours                                                                                                              |
| Result:          | Inhibited cell growth of RPMI8226 and U87MG cells with IC <sub>50</sub> of 1.6 $\mu$ M and 2.4 $\mu$ M, respectively. |

#### Western Blot Analysis<sup>[1]</sup>

|                  |                                                                       |
|------------------|-----------------------------------------------------------------------|
| Cell Line:       | HEK293T cells                                                         |
| Concentration:   | 0 $\mu$ M, 3 $\mu$ M, 10 $\mu$ M                                      |
| Incubation Time: | 24 hours                                                              |
| Result:          | Enhanced proteolytic degradation in a concentration-dependent manner. |

## CUSTOMER VALIDATION

- Eur J Pharmacol. 2023 Aug 24;176011.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Njomen E, et al. Small Molecule Modulation of Proteasome Assembly. Biochemistry. 2018 Jul 17;57(28):4214-4224.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA