



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

(2S,3R)-Brassinazole

Cat. No.:	HY-121161B		
CAS No.:	259200-31-2		
Molecular Formula:	C ₁₈ H ₁₈ ClN ₃ O		
Molecular Weight:	327.81		
Target:	Drug Isomer		
Pathway:	Others		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 20 mg/mL (61.01 mM; ultrasonic and warming and heat to 60°C)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		3.0505 mL	15.2527 mL	30.5055 mL
	5 mM		0.6101 mL	3.0505 mL	6.1011 mL
	10 mM		0.3051 mL	1.5253 mL	3.0505 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

(2S,3R)-Brassinazole, the enantiomer of Brassinazole (BRZ). Brassinazole inhibits brassinosteroid (BR) biosynthesis, via acting on the oxidative processes from 6-oxo-campestanol to teasterone. (2S,3R)-Brassinazole might be the most active form of Brz^{[1][2][3]}.

IC₅₀ & Target

Brassinosteroid (BR) biosynthesis^[1]

In Vitro

Brassinazole (0.5, 1, 5 μM) causes markedly deformed seedlings, whose morphology is similar to that of BR-deficient mutants. Brassinazole causes cress dwarfism, altering leaf morphology such as the typical downward curl and dark green appearance of Arabidopsis BR-deficient mutants. However, administration of 10 nM BR reversed dwarfism^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Yoshida Shigeo, et al. Preparation of (1,2,4-triazolyl)alkanols as specific inhibitors of brassinosteroid biosynthesis: Japan, JP2000053657. 2000-02-22.

[2]. Asami T, et al. Mode of action of brassinazole: a specific inhibitor of brassinosteroid biosynthesis[M]. 2000.

[3]. T Asami, et al. Characterization of Brassinazole, a Triazole-Type Brassinosteroid Biosynthesis Inhibitor. Plant Physiol. 2000 May;123(1):93-100.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA