



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)


[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



## IM-12

Cat. No.:	HY-12292
CAS No.:	1129669-05-1
Molecular Formula:	C <sub>22</sub> H <sub>20</sub> FN <sub>3</sub> O <sub>2</sub>
Molecular Weight:	377
Target:	GSK-3
Pathway:	PI3K/Akt/mTOR; Stem Cell/Wnt
Storage:	<div> Powder -20°C 3 years </div> <div> 4°C 2 years </div> <div> In solvent -80°C 2 years </div> <div> -20°C 1 year </div>



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : ≥ 54.9 mg/mL (145.62 mM)  
 \* "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.6525 mL	13.2626 mL	26.5252 mL
	5 mM		0.5305 mL	2.6525 mL	5.3050 mL
	10 mM		0.2653 mL	1.3263 mL	2.6525 mL

Please refer to the solubility information to select the appropriate solvent.

#### In Vivo

1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
 Solubility: ≥ 2.5 mg/mL (6.63 mM); Clear solution

### BIOLOGICAL ACTIVITY

Description	IM-12 is an inhibitor of GSK-3β, with an IC <sub>50</sub> of 53 nM, and also enhances Wnt signalling.
IC <sub>50</sub> & Target	GSK-3β 53 nM (IC <sub>50</sub> )
In Vitro	<p>IM-12 inhibits GSK-3β in ReNcell VM cells, with I<sub>50</sub> of 3.8 μM. IM-12 (3 μM) enhances the β-catenin amount, with no further effect at lower or higher concentration. IM-12 (3 μM) also attenuates the proliferation of ReNCell VM cells. IM-12 increases TCF-activity of ReNcell VM<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

## PROTOCOL

### Kinase Assay <sup>[1]</sup>

Cells are lysed in RIPA buffer, supplemented with protease and phosphatase inhibitors and centrifuged for 5 min at 15,000 rpm. Immunoprecipitation of GSK-3 $\beta$  is performed with a specific mouse monoclonal anti GSK-3 $\beta$  [G8] antibody with 5  $\mu$ g/sample for 2 h at 4°C. The bound protein is precipitated with Protein A/G-Plus agarose-beads (10  $\mu$ L beads per sample). GSK-3 $\beta$  kinase activity is measured in a reaction mixture containing final concentrations of: 4 mM MOPS pH 7.2; 0.4 mM EDTA; 1 mM EGTA; 2.5 mM  $\beta$ -glycerophosphate; 4 mM MgCl<sub>2</sub>; 40  $\mu$ M BSA; 0.05 mM DTT. 10  $\mu$ g/sample pGS-2 peptide substrate is used<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### Cell Assay <sup>[1]</sup>

To measure viable cells, 50-100  $\mu$ L of cell suspension is analyzed using CASY technology with the appropriate program. ReNcell VM cells are seeded at a defined cell number and proliferated for 24 h. Then the medium is changed to proliferation medium with added substances at indicated concentrations. The cell number is determined every 24 h. Cells are exposed to the added drugs during the whole experiment, whereas the media is changed every 24 h<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- J Cell Physiol. 2019 Jul;234(7):10698-10708.
- Carcinogenesis. 2023 Jan 16;bgad003.
- Mol Med Rep. 2022 Oct;26(4):306.
- Methods Mol Biol. 2023 Jun 24.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Schmole AC, et al. Novel indolylmaleimide acts as GSK-3 $\beta$  inhibitor in human neural progenitor cells. Bioorg Med Chem. 2010 Sep 15;18(18):6785-95.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA