



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Cystamine

Cat. No.:	HY-124476
CAS No.:	51-85-4
Molecular Formula:	C ₄ H ₁₂ N ₂ S ₂
Molecular Weight:	152.28
Target:	Caspase; Glutaminase; Apoptosis
Pathway:	Apoptosis; Metabolic Enzyme/Protease
Storage:	Pure form -20°C 3 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (656.69 mM; Need ultrasonic)					
	H ₂ O : 100 mg/mL (656.69 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		6.5669 mL	32.8343 mL	65.6685 mL
		5 mM		1.3134 mL	6.5669 mL	13.1337 mL
		10 mM		0.6567 mL	3.2834 mL	6.5669 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (16.42 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (16.42 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (16.42 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	Cystamine is the disulfide form of the free thiol, cysteamine. Cystamine is an orally active transglutaminase (Tgase) inhibitor. Cystamine also has inhibition activity for caspase-3 with an IC ₅₀ value of 23.6 μM. Cystamine can be used for the research of severals diseases including Huntington's disease (HD) ^{[1][2][3]} .
IC ₅₀ & Target	Caspase 3 23.6 μM (IC ₅₀)

In Vitro

Cystamine has inhibition activity for caspase-3 with an IC_{50} value of 23.6 μM ^[1].
Cystamine (0-500 μM ; 0-16 h) inhibits recombinant active caspase-3 in a concentration-dependent manner^[1].
Cystamine (250 μM ; 10 h) robustly increases the levels of glutathione^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.
Western Blot Analysis^[1]

Cell Line:	uman neuroblastoma SH-SY5Y cells
Concentration:	250, 500 μM
Incubation Time:	0-16 h
Result:	Inhibited the MG132-mediated activation of caspase-3. Inhibited the H2O2-mediated activation of caspase-3. Inhibited caspase-3 activity in a tTG-independent manner.

In Vivo

Cystamine (oral, i.p.; 112, 225 mg/kg) reduces Tgase activity and GGEL levels, lessens the behavioral and neuropathological severity, and extends survival in R6/2 transgenic HD mice^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	R6/2 transgenic HD mice ^[2]
Dosage:	112, 225 mg/kg
Administration:	Intraperitoneal or oral, daily
Result:	Significantly extended survival, improved body weight and motor performance, delayed the neuropathological sequela and significantly altered the levels of Tgase activity and N(Sigma)-(gamma-L-glutamyl)-L-lysine (GGEL) levels.

REFERENCES

- [1]. Mathieu Lesort, et al. Cystamine inhibits caspase activity. Implications for the treatment of polyglutamine disorders. J Biol Chem. 2003 Feb 7;278(6):3825-30.
- [2]. Alpaslan Dedeoglu, et al. Therapeutic effects of cystamine in a murine model of Huntington's disease. J Neurosci. 2002 Oct 15;22(20):8942-50.
- [3]. Thomas M Jeitner, et al. Cystamine and cysteamine as inhibitors of transglutaminase activity in vivo. Biosci Rep. 2018 Sep 5;38(5):BSR20180691.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA