



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Ro-3306

Cat. No.:	HY-12529
CAS No.:	872573-93-8
Molecular Formula:	C ₁₈ H ₁₃ N ₃ OS ₂
Molecular Weight:	351
Target:	CDK; Apoptosis
Pathway:	Cell Cycle/DNA Damage; Apoptosis
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 1 year -20°C 6 months



SOLVENT & SOLUBILITY

In Vitro	DMSO : 25 mg/mL (71.23 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		2.8490 mL	14.2450 mL	28.4900 mL
		5 mM		0.5698 mL	2.8490 mL	5.6980 mL
		10 mM		0.2849 mL	1.4245 mL	2.8490 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 0.5% CMC-Na/saline water Solubility: 5 mg/mL (14.25 mM); Suspended solution; Need ultrasonic					
	2. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 1.67 mg/mL (4.76 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 1.67 mg/mL (4.76 mM); Suspended solution					

BIOLOGICAL ACTIVITY

Description	Ro-3306 is a potent and selective inhibitor of CDK1, with K _i s of 20 nM, 35 nM and 340 nM for CDK1, CDK1/cyclin B1 and CDK2/cyclin E, respectively.			
IC ₅₀ & Target	CDK1 20 nM (Ki)	CDK1/cyclinB1 35 nM (Ki)	CDK1/cyclin A 110 nM (Ki)	CDK2/cyclinE 340 nM (Ki)
	PKCδ 318 nM (Ki)	SGK 497 nM (Ki)	ERK 1980 nM (Ki)	

In Vitro

RO-3306 is an ATP-competitive inhibitor, and inhibits CDK1/cyclin A complexes with K_i of 110 nM. RO-3306 blocks the cell cycle in the G2/M phase of human cancer cells. RO-3306 (4 μ M) induces apoptosis in cancer cells^[1]. RO-3306 (5 μ M) induces G2/M-phase cell cycle arrest and apoptosis of AML cells in a time-dependent manner. RO-3306 treatment significantly increases the percentage of Annexin V-positive cells in G1-phase cells without affecting the cell cycle distribution. RO-3306 enhances p53-mediated apoptosis. RO-3306 cooperates with Nutlin-3 in activating Bax and inducing mitochondrial apoptosis. RO-3306 (5 μ M) downregulates antiapoptotic p21, Bcl-2 and survivin protein expression in AML. RO-3306 inhibits p53-induced p21 synthesis. RO-3306 does not inhibit RNA polymerase II CTD phosphorylation^[2]. RO-3306 (10 μ M) effectively arrests oocyte maturation. RO-3306 reduces the blastocyst formation in oocytes^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Kinase Assay ^[1]

The CDK assays are run by using recombinant human CDK/cyclin complexes (CDK1/cyclin B1, CDK1/cyclin A, CDK2/cyclin E, and CDK4/cyclin D) expressed and isolated from Hi5 insect cells. GST-cyclin B1, CDK1, GST-cyclin-E, CDK2, GST-CDK4, and cyclin D, are used in the assay. The GST-tagged proteins are coexpressed and purified in complex with their partners. All assays use a His-6-tagged fragment of pRB (amino acids 385-928) as a substrate. The protein is expressed from a construct. It is expressed in M15 Escherichia coli cells and bound on a Ni-chelated agarose column pretreated with 1 mM imidazole and eluted with 500 mM imidazole. The eluted protein is dialyzed against 20 mM Hepes, pH 7/6.25 mM $MgCl_2$ /1.5 mM DTT, aliquoted, and stored at $-80^\circ C$.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Nat Commun. 2023 May 26;14(1):3050.
- Nat Commun. 2023 Apr 6;14(1):1919.
- Cancer Commun (Lond). 2021 Jan 20.
- Nucleic Acids Res. 2023 Jan 18;gkac1269.
- Cell Death Differ. 2021 Feb;28(2):799-813.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Vassilev LT, et al. Selective small-molecule inhibitor reveals critical mitotic functions of human CDK1. Proc Natl Acad Sci U S A. 2006 Jul 11;103(28):10660-5.
- [2]. Kojima K, et al. Cyclin-dependent kinase 1 inhibitor RO-3306 enhances p53-mediated Bax activation and mitochondrial apoptosis in AML. Cancer Sci. 2009 Jun;100(6):1128-36.
- [3]. Jang WI, et al. A specific inhibitor of CDK1, RO-3306, reversibly arrests meiosis during in vitro maturation of porcine oocytes. Anim Reprod Sci. 2014 Jan 30;144(3-4):102-8.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA