



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## IWP-4

Cat. No.:	HY-12879		
CAS No.:	686772-17-8		
Molecular Formula:	C <sub>23</sub> H <sub>20</sub> N <sub>4</sub> O <sub>3</sub> S <sub>3</sub>		
Molecular Weight:	496.62		
Target:	Wnt		
Pathway:	Stem Cell/Wnt		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



## SOLVENT & SOLUBILITY

### In Vitro

DMSO : 1.3 mg/mL (2.62 mM; Need ultrasonic)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.0136 mL	10.0681 mL	20.1361 mL
	5 mM		---	---	---
	10 mM		---	---	---

Please refer to the solubility information to select the appropriate solvent.

## BIOLOGICAL ACTIVITY

Description	IWP-4 is a small molecule Wnt inhibitor with an IC <sub>50</sub> of 25 nM.
IC <sub>50</sub> & Target	IC <sub>50</sub> : 25 nM (Wnt) <sup>[1]</sup>
In Vitro	<p>IWP-4 is a small molecule Wnt inhibitor with an IC<sub>50</sub> of 25 nM. IWP-4 induces the expression of cardiac markers, including cardiac troponin I (CTNI) and cardiac myosin heavy chain bright cells (MYH<sup>hi+</sup>). IWP-4 also results in the appearance of beating foci (0.44±0.10 SEM beats per second), which is absent in all cultures not receiving IWP-4. Further, flow cytometric analysis shows that there are significantly more MYH<sup>lo+</sup> cells in IWP-4 treated cultures (P&lt;0.0002) compare with untreated cultures at day 16, being 17.0±1.3 SD% and 5.4±1.4 SD%, respectively. Quantification of NKX2-5 protein expression shows that 63% (481/817) of IWP-4 treated cells display nuclear NKX2-5 expression<sup>[1]</sup>. Mesenchymal precursor cells (MPCs) treated with IWP-4 show no significant changes in the expression of AXIN2, CTNNB1 and GSK3B as compare to osteogenic medium alone on day 7, but MPCs treated with IWP-4 express elevates levels of DKK1 and GSK3β on day 21. IWP-4 also causes a significant down regulation of SPARC and COL1A1<sup>[2]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

---

## PROTOCOL

### Cell Assay <sup>[1]</sup>

hESC cultures are obtained in mTeSR-1 medium and expanded with daily medium exchange until colonies reach the desired level of confluence (~70% to 80%). At this time (marked day 0), mTeSR-1 is replaced with a basal medium comprised of RPMI 1640 medium supplemented with 2% B27 supplement and 1% penicillin/streptomycin. 20 ng/mL BMP-4 and/or 6 ng/mL activin A are added to the basal medium for primitive streak induction, and exchanged daily until day 3. Then, basal media with or without 5 mM IWP-4 is added to the cells and exchanged every 2 days [dimethyl sulfoxide (DMSO) at the same concentration is used as a vehicle control] until day 15, after which basal medium is supplied every 2 days<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Chin Med. 2022 Jan 6;17(1):11.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Hudson J, et al. Primitive cardiac cells from human embryonic stem cells. Stem Cells Dev. 2012 Jun 10;21(9):1513-23.

[2]. Frith JE, et al. Microbioreactor array screening of Wnt modulators and microenvironmental factors in osteogenic differentiation of mesenchymal progenitor cells. PLoS One. 2013 Dec 23;8(12):e82931.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA