



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Clomethiazole

Cat. No.:	HY-129105
CAS No.:	533-45-9
Molecular Formula:	C ₆ H ₈ ClNS
Molecular Weight:	161.65
Target:	GABA Receptor; Cytochrome P450
Pathway:	Membrane Transporter/Ion Channel; Neuronal Signaling; Metabolic Enzyme/Protease
Storage:	Pure form -20°C 3 years 4°C 2 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (618.62 mM; Need ultrasonic)				
	Preparing Stock Solutions	Solvent Concentration Mass	1 mg	5 mg	10 mg
		1 mM	6.1862 mL	30.9310 mL	61.8621 mL
		5 mM	1.2372 mL	6.1862 mL	12.3724 mL
		10 mM	0.6186 mL	3.0931 mL	6.1862 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (12.87 mM); Clear solution				
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (12.87 mM); Clear solution				
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (12.87 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	Chlormethiazole is a potent and orally active GABA _A agonist ^[1] . Chlormethiazole inhibits cytochrome P450 isoforms: CYP2A6 and CYP2E1 in human liver microsomes. Chlormethiazole is an anticonvulsant agent and has the potential for treating convulsive status epilepticus ^[2] .
IC ₅₀ & Target	CYP2
In Vitro	Clomethiazole (1 mM), in the absence of GABA, to α1/β1/γ2 or α1/β2/γ2 subunit-containing cells produced large whole-cell

currents^[1].

Clomethiazole activate GABAA currents in $\alpha 1/\beta 1/\gamma 2$ - and $\alpha 1/\beta 2/\gamma 2$ -containing cells, with EC₅₀ values of 0.3 and 1.5 mM, respectively^[1].

Clomethiazole (30 μ M) at low concentration also potentiates the action of GABA in both cell types, equivalent to a 3-fold increase in potency and up to 1.8-fold increase in maximal current^[1].

Clomethiazole inhibits cytochrome P450 isoforms, CYP2A6 and CYP2E1 with IC₅₀ values of 24 μ M and 42 μ M, respectively, in human liver microsomes, meanwhile all other isoforms exhibiting values > 300 μ M^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Biomaterials. 4 August 2022, 121720.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Nelson RM, et al. Electrophysiological actions of gamma-aminobutyric acid and clomethiazole on recombinant GABA(A) receptors. Eur J Pharmacol. 2002 Oct 11;452(3):255-62.

[2]. Stresser DM, et al. Selective Time- and NADPH-Dependent Inhibition of Human CYP2E1 by Clomethiazole. Drug Metab Dispos. 2016 Aug;44(8):1424-30

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA