



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Stearoyl-L-carnitine-d₃ chloride

Cat. No.:	HY-130466S
CAS No.:	2245711-27-5
Molecular Formula:	C ₂₅ H ₄₇ D ₃ ClNO ₄
Molecular Weight:	467.14
Target:	GlyT; Endogenous Metabolite
Pathway:	Membrane Transporter/Ion Channel; Neuronal Signaling; Metabolic Enzyme/Protease
Storage:	4°C, sealed storage, away from moisture * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro

Ethanol : ≥ 20 mg/mL (42.81 mM)
 Ethanol : ≥ 20 mg/mL (42.81 mM)
 DMF : ≥ 20 mg/mL (42.81 mM)
 DMF : ≥ 20 mg/mL (42.81 mM)
 DMSO : ≥ 14 mg/mL (29.97 mM)
 * "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.1407 mL	10.7034 mL	21.4069 mL
	5 mM		0.4281 mL	2.1407 mL	4.2814 mL
	10 mM		0.2141 mL	1.0703 mL	2.1407 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

Stearoyl-L-carnitine-d₃ (chloride) is the deuterium labeled Stearoyl-L-carnitine chloride. Stearoyl-L-carnitine chloride is an endogenous long-chain acylcarnitine. Stearoyl-L-carnitine chloride is a less potent inhibitor of GlyT2. Stearoyl-L-carnitine chloride inhibits glycine responses by 16.8% at concentrations up 3 μM^{[1][2]}.

IC₅₀ & Target

GlyT2

In Vitro

Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs^[1].
 MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother*. 2019;53(2):211-216.
 - [2]. Cristofano A, et, al. Serum Levels of Acyl-Carnitines along the Continuum from Normal to Alzheimer's Dementia. *PLoS One*. 2016 May 19;11(5):e0155694.
 - [3]. Carland JE, et, al. Oleoyl-L-carnitine inhibits glycine transport by GlyT2. *Br J Pharmacol*. 2013 Feb;168(4):891-902.
 - [4]. Huang W, et, al. Carnitine transport and its inhibition by sulfonylureas in human kidney proximal tubular epithelial cells. *Biochem Pharmacol*. 1999 Oct 15;58(8):1361-70.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA