

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Proteins



MK-8245

Cat. No.: HY-13070 CAS No.: 1030612-90-8 Molecular Formula: C₁₇H₁₆BrFN₆O₄

Molecular Weight: 467.25

Target: Stearoyl-CoA Desaturase (SCD) Pathway: Metabolic Enzyme/Protease Storage: 4°C, protect from light

* In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)

Product Data Sheet

SOLVENT & SOLUBILITY

DMSO : ≥ 100 mg/mL (214.02 mM) In Vitro

* "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	2.1402 mL	10.7009 mL	21.4018 mL
	5 mM	0.4280 mL	2.1402 mL	4.2804 mL
	10 mM	0.2140 mL	1.0701 mL	2.1402 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (5.35 mM); Clear solution
- 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (5.35 mM); Clear solution
- 3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (5.35 mM); Clear solution

BIOLOGICAL ACTIVITY

Description	MK-8245 is a potent, liver-targeted stearoyl-CoA desaturase (SCD) inhibitor, with IC $_{50}$ s of 1 nM for human SCD1 and 3 nM for both rat SCD1 and mouse SCD1, with antidiabetic and antidyslipidemic efficacy $^{[1]}$.	
IC ₅₀ & Target	IC50: 1 nM (human SCD1), 3 nM (rat SCD1), 3 nM (mouse SCD1) ^[1]	
In Vitro	MK-8245 is a potent and liver-specific SCD inhibitor ^[1] . MK-8245 displays similar potencies against human, rat and mouse SCD1, with IC ₅₀ values of 1 nM for human SCD1 and 3 nM for both rat SCD1 and mouse SCD1 ^[1] .	

MK-8245 exhibits a significant SCD inhibition in the rat hepatocyte assay which contains functional, actives organic anion transporting polypeptides (OATPs) with IC₅₀ of 68 nM, while being only weakly active OATPs in the HepG2 cell assay which is devoid of active with IC₅₀ of $^{-1} \mu M^{[1]}$.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

MK-8245 (10mg/kg; p.o.) exhibits a tissue distribution profile concentrated in the liver, with low exposure in tissues associated with potential adverse events in rats, dogs, and rhesus monkeys^[1].

MK-8245 improves glucose clearance in a dose-dependent manner in eDIO mice administrated before the glucose challenge [1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Male C57BL6 mice, male Sprague-Dawley rats ^[1]	
Dosage:	10mg/kg	
Administration:	Oral administration	
Result:	Exhibits a tissue distribution profile concentrated in the liver.	

CUSTOMER VALIDATION

- Sci China Life Sci. 2021 May 27;1-21.
- Aquaculture. 2023 Jun 10, 739766.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Oballa RM, et al. Development of a liver-targeted stearoyl-CoA desaturase (SCD) inhibitor (MK-8245) to establish a therapeutic window for the treatment of diabetes and dyslipidemia. J Med Chem. 2011 Jul 28;54(14):5082-96. Epub 2011 Jun 28.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA