



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

BOC-D-FMK

Cat. No.:	HY-13229
CAS No.:	634911-80-1
Molecular Formula:	C ₁₁ H ₁₈ FNO ₅
Molecular Weight:	263.26
Target:	Caspase; Apoptosis
Pathway:	Apoptosis
Storage:	Pure form -20°C 3 years 4°C 2 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (379.85 mM; Need ultrasonic)				
		Mass			
		Solvent	1 mg	5 mg	10 mg
		Concentration			
	Preparing Stock Solutions	1 mM	3.7985 mL	18.9926 mL	37.9853 mL
		5 mM	0.7597 mL	3.7985 mL	7.5971 mL
		10 mM	0.3799 mL	1.8993 mL	3.7985 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (9.50 mM); Clear solution 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (9.50 mM); Clear solution 3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (9.50 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	Boc-D-FMK is a cell-permeable, irreversible and broad spectrum caspase inhibitor. Boc-D-FMK inhibits apoptosis stimulated by TNF-α with an IC ₅₀ of 39 μM.
IC ₅₀ & Target	Caspase
In Vitro	Apoptosis is a pathway of cell death orchestrated by a family of proteases called caspases. Boc-D-fmk inhibits TNFα-stimulated reactive oxygen species (ROS) generation. Boc-D-FMK inhibits apoptosis stimulated by TNF-α with an IC ₅₀ of 39

μM ^[1]. BocD-fmk at 50 μM prevents genistein-induced apoptosis of p815 cells. Confocal microscopy shows that the release of mitochondrial apoptotic factors is inhibited by BocD-fmk^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Boc-D-FMK-fmk effectively attenuates the hepatocyte apoptosis in bile duct-ligated rats and may improve the survival rates after endotoxin challenge^[3]. A single injection of Boc-D-FMK results in longterm protection of MNs against root avulsion-induced death for more than 8 weeks and the Boc-D-FMK-treated MNs are able to regenerate their axons into an implanted PN graft and reinnervate the target muscle^[4].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Animal Administration ^[3]

Rats: Boc-D-FMK is dissolved in DMSO. Male Sprague-Dawley rats group 1 (OBBOC-D) undergo common bile duct ligation and simultaneously treatment with Boc-D-FMK. The first dose of Boc-D-FMK (1.5 mg/kg) is injected into the inferior vena cava immediately after bile duct ligation. Subsequent doses of Boc-DFMK (1.5 mg/kg twice daily) are given intraperitoneally on the first and second postoperative days. The last dose (1.5 mg/kg) is given on the morning of the third postoperative day ^[3].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- J Exp Clin Cancer Res. 2023 Jun 6;42(1):142.
- Anal Chem. 2017 Sep 19;89(18):9788-9796.
- Sci Rep. 2017 Jun 7;7(1):2929.
- Saudi J Biol Sci. 6 December 2021.
- Exp Cell Res. 2018 Sep 1;370(1):103-115.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Cowburn AS, et al. z-VAD-fmk augmentation of TNF alpha-stimulated neutrophil apoptosis is compound specific and does not involve the generation of reactive oxygen species.
- [2]. Yee SB, et al. zVAD-fmk, unlike BocD-fmk, does not inhibit caspase-6 acting on 14-3-3/Bad pathway in apoptosis of p815 mastocytoma cells. Exp Mol Med. 2006 Dec 31;38(6):634-42.
- [3]. Sheen-Chen SM, et al. Effect of Boc-D-Fmk on hepatocyte apoptosis after bile duct ligation in rat and survival rate after endotoxin challenge. J Gastroenterol Hepatol. 2008 Aug;23(8 Pt 1):1276-9.
- [4]. Chan YM, et al. Inhibition of caspases promotes long-term survival and reinnervation by axotomized spinal motoneurons of denervated muscle in newborn rats. Exp Neurol. 2003 Jun;181(2):190-203.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA