



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Efaproxiral

Cat. No.:	HY-13619
CAS No.:	131179-95-8
Molecular Formula:	C ₂₀ H ₂₃ NO ₄
Molecular Weight:	341.4
Target:	Reactive Oxygen Species
Pathway:	Immunology/Inflammation; Metabolic Enzyme/Protease; NF-κB
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 150 mg/mL (439.37 mM)

* "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.9291 mL	14.6456 mL	29.2912 mL
	5 mM		0.5858 mL	2.9291 mL	5.8582 mL
	10 mM		0.2929 mL	1.4646 mL	2.9291 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 50% PEG300 >> 50% saline
Solubility: 27.5 mg/mL (80.55 mM); Clear solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.08 mg/mL (6.09 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.08 mg/mL (6.09 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.08 mg/mL (6.09 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Efaproxiral is a haemoglobin (Hb) synthetic allosteric modifier, decreases Hb-oxygen (O₂) binding affinity and enhances oxygenation of hypoxic tumours during radiation therapy ^[1].

IC₅₀ & Target

haemoglobin (Hb)^[1]

In Vitro	Efaproxiral binds to only one pair of symmetry-related sites in the Hb central water cavity^[2]. Efaproxiral readily crosses the red cell membrane in the presence of serum albumin solutions^[2]. Efaproxiral is not inhibited from entering erythrocytes in the presence of an anion-channel blocking agent (DIDS)^[2]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.								
In Vivo	Efaproxiral (150 mg/kg, i.p.) increase tumor oxygenation and increase the tumor growth inhibition of radiotherapy over 5 days of treatment^[3]. Efaproxiral reduces hemoglobin-oxygen binding affinity, which facilitates oxygen release from hemoglobin into surrounding tissues and potentially increases the pO₂ of the tumors^[4] MCE has not independently confirmed the accuracy of these methods. They are for reference only. <table> <tr> <td>Animal Model:</td><td>Female C3H/HEJ mice (18–20 g), with radiation-induced fibrosarcoma tumor (RIF-1) cells xenograft^[3]</td></tr> <tr> <td>Dosage:</td><td>150 mg/kg</td></tr> <tr> <td>Administration:</td><td>Intraperitoneal injection; prior to X Irradiation (4 Gy/day), for 5 days</td></tr> <tr> <td>Result:</td><td>Significantly increased tumor oxygenation by 8.4 to 43.4 mmHg within 5 days, with maximum increases at 22–31 minutes after treatment.</td></tr> </table>	Animal Model:	Female C3H/HEJ mice (18–20 g), with radiation-induced fibrosarcoma tumor (RIF-1) cells xenograft ^[3]	Dosage:	150 mg/kg	Administration:	Intraperitoneal injection; prior to X Irradiation (4 Gy/day), for 5 days	Result:	Significantly increased tumor oxygenation by 8.4 to 43.4 mmHg within 5 days, with maximum increases at 22–31 minutes after treatment.
Animal Model:	Female C3H/HEJ mice (18–20 g), with radiation-induced fibrosarcoma tumor (RIF-1) cells xenograft ^[3]								
Dosage:	150 mg/kg								
Administration:	Intraperitoneal injection; prior to X Irradiation (4 Gy/day), for 5 days								
Result:	Significantly increased tumor oxygenation by 8.4 to 43.4 mmHg within 5 days, with maximum increases at 22–31 minutes after treatment.								

CUSTOMER VALIDATION

- J Enzyme Inhib Med Chem. 2021 Dec;36(1):377-383.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Stea B, et al. Efaproxiral red blood cell concentration predicts efficacy in patients with brain metastases. Br J Cancer. 2006 Jun 19;94(12):1777-1784.
- [2]. Abraham DJ, et al. Allosteric modifiers of hemoglobin: 2-[4-[(3,5-disubstituted anilino)carbonyl]methyl]phenoxy]-2-methylpropionic acid derivatives that lower the oxygen affinity of hemoglobin in red cell suspensions, in whole blood, and in vivo in rats.
- [3]. Hou H, et al. The effects of Efaproxyn (efaproxiral) on subcutaneous RIF-1 tumor oxygenation and enhancement of radiotherapy-mediated inhibition of tumor growth in mice. Radiat Res. 2007 Aug;168(2):218-25.
- [4]. Hou H, et al. Increased oxygenation of intracranial tumors by efaproxyn (efaproxiral), an allosteric hemoglobin modifier: In vivo EPR oximetry study. Int J Radiat Oncol Biol Phys. 2005 Apr 1;61(5):1503-9.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA