

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Proteins

Product Data Sheet

Entecavir monohydrate

Cat. No.: HY-13623A CAS No.: 209216-23-9 Molecular Formula: C₁₂H₁₇N₅O₄ Molecular Weight: 295.29 HBV Target:

Pathway: Anti-infection

Storage: Powder -20°C 3 years

2 years

-80°C In solvent 2 years

> -20°C 1 year

SOLVENT & SOLUBILITY

In Vitro

DMSO: $\geq 50 \text{ mg/mL} (169.33 \text{ mM})$

H₂O: 2.8 mg/mL (9.48 mM; Need ultrasonic and warming)

* "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	3.3865 mL	16.9325 mL	33.8650 mL
	5 mM	0.6773 mL	3.3865 mL	6.7730 mL
	10 mM	0.3387 mL	1.6933 mL	3.3865 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 3 mg/mL (10.16 mM); Clear solution
- 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 3 mg/mL (10.16 mM); Clear solution
- 3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 3 mg/mL (10.16 mM); Clear solution

BIOLOGICAL ACTIVITY

Description Entecavir monohydrate (BMS200475 monohydrate; SQ34676 monohydrate) is a potent and selective inhibitor of HBV, with an EC₅₀ of 3.75 nM in HepG2 cell.

EC50\(\text{\text{\text{3}}}\).75 nM (anti-HBV, HepG2 cell)\([1]) IC₅₀ & Target

In Vitro	Entecavir monohydrate (BMS200475 monohydrate; SQ34676 monohydrate) has a EC $_{50}$ of 3.75 nM against HBV. It is incorporated into the protein primer of HBV and subsequently inhibits the priming step of the reverse transcriptase. The antiviral activity of BMS-200475 is significantly less against the other RNA and DNA viruses ^[1] . Entecavir monohydrate is more readily phosphorylated to its active metabolites than other deoxyguanosine analogs (penciclovir, ganciclovir, lobucavir, and aciclovir) or lamivudine. The intracellular half-life of entecavir is 15 h ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	Daily oral treatment with Entecavir monohydrate at doses ranging from 0.02 to 0.5 mg/kg of body weight for 1 to 3 months effectively reduces the level of woodchuck hepatitis virus (WHV) viremia in chronically infected woodchucks ^[3] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay [1]

BMS 200475 is prepared in phosphate-buffered saline (PBS) and diluted with appropriate medium containing 2% fetal bovine serum. HepG2 2.2.15 cells are plated at a density of 5×10^5 cells per well on 12-well Biocoat collagen-coated plates and are maintained in a confluent state for 2 to 3 days before being overlaid with 1 mL of medium spiked with BMS 200475. Quantification of HBV was performed on day $10^{[1]}$.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Adv Sci (Weinh). 2022 May;9(16):e2103135.
- Emerg Microbes Infect. 2020 Dec 9;1-22.
- Antiviral Res. 2020 Aug;180:104826.
- Cell Mol Gastroenterol Hepatol. 2021 Dec 8;S2352-345X(21)00249-6.
- PLoS Pathog. 2021 Aug 9;17(8):e1009838.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Innaimo SF, et al. Identification of?BMS-200475?as a potent and selective inhibitor of hepatitis B virus. Antimicrob Agents Chemother. 1997 Jul;41(7):1444-8.

[2]. Rivkin A, et al. A review of entecavir in the treatment of chronic hepatitis B infection. Curr Med Res Opin. ?2005 Nov;21(11):1845-56.

[3]. Genovesi EV, et al. Efficacy of the carbocyclic 2'-deoxyguanosine nucleoside?BMS-200475?in the woodchuck model of hepatitis B virus infection. Antimicrob Agents Chemother.?1998 Dec;42(12):3209-17.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA