



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## JIB-04

Cat. No.:	HY-13953
CAS No.:	199596-05-9
Molecular Formula:	C <sub>17</sub> H <sub>13</sub> ClN <sub>4</sub>
Molecular Weight:	308.76
Target:	Histone Demethylase; Apoptosis
Pathway:	Epigenetics; Apoptosis
Storage:	Powder    -20°C    3 years 4°C    2 years In solvent   -80°C    2 years -20°C    1 year



### SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (161.94 mM; Need ultrasonic)					
	Ethanol : 2 mg/mL (6.48 mM; Need ultrasonic)					
	Preparing Stock Solutions	<div><div>Solvent</div><div>Concentration</div></div>	Mass	1 mg	5 mg	10 mg
		1 mM		3.2387 mL	16.1935 mL	32.3871 mL
		5 mM		0.6477 mL	3.2387 mL	6.4774 mL
		10 mM		0.3239 mL	1.6194 mL	3.2387 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 0.5% CMC-Na/saline water Solubility: 10 mg/mL (32.39 mM); Suspended solution; Need ultrasonic					
	2. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (6.74 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 1.25 mg/mL (4.05 mM); Clear solution					

### BIOLOGICAL ACTIVITY

Description	JIB-04 is a pan-selective Jumonji histone demethylase inhibitor with IC <sub>50</sub> s of 230, 340, 855, 445, 435, 1100, and 290 nM for JARID1A, JMJD2E, JMJD3, JMJD2A, JMJD2B, JMJD2C, and JMJD2D, respectively.		
IC <sub>50</sub> & Target	KDM4	KDM6	KDM5
In Vitro	JIB-04 is consistently selective for cancer vs. normal cells, demonstrated by the higher sensitivity of lung and prostate		

cancer lines (with IC<sub>50</sub> as low as 10 nM) compared to HBECs and PrSCs/PrECs. JIB-04 inhibits cellular Jumonji demethylase activity, and Jumonji levels affect JIB-04 action in cells<sup>[1]</sup>. JIB-04 significantly inhibits the proliferation of GB cell lines and stem-enriched cultures. JIB-04 exerts its maximal inhibitory activity against KDM5A, and modulates the expression of genes involved in the control of cancer cell growth and leads to hypermethylation of H3K4. Furthermore, JIB-04 (2500 nM) activates the autophagy and apoptotic pathways and inactivates PI3K. JIB-04 also cooperates with TMZ in killing GB cells<sup>[2]</sup>. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### In Vivo

JIB-04 results in a significant reduction in cancer-induced death rates in mice, prolonging survival<sup>[1]</sup>. JIB-04 (60, 40 and 20 mg/kg, i.p.) reaches bioactive concentration in the brain of the mice. The orthotopic GB xenograft model shows a trend toward longer survival in JIB-04-treated mice with an Hazard Ratio of 0.5<sup>[2]</sup>. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

### Cell Assay <sup>[1]</sup>

For cell viability assays, cells are plated at 1500-3000 cells/well in 96 well plates and treated the next day with increasing doses of compound over 4 days and their viability assessed by standard MTS assays using Promega's Cell Titer or Cell Titer-Glo reagents. Absorbance at 490 nm and 650 nm or luminescence is measured by a Spectra Max or a FlouoroStar Omega plate reader. Data are normalized to the untreated controls (100% viability). Each cell line is tested in 2-5 independent assays, each containing 4-8 replicates. IC<sub>50</sub> values are calculated using DIVISA, a high-throughput software, developed in house, for storing and analyzing drug sensitivity assays. Dose-response curves are plotted using a non-linear regression model and IC<sub>50</sub>s are determined from the fitted curves. The average IC<sub>50</sub> derived from 2-5 independent assays, each containing 4-8 replicates is reported.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### Animal Administration <sup>[1]</sup>

For xenografts, 4-6 week old female nude mice are housed under standard conditions in a clean facility at UTSW. Two million H358 cells or five million A549 cells are injected subcutaneously and allowed to grow for 2-3 weeks with monitoring. When tumors reach appr 200 mm<sup>3</sup>, therapy is started in weight and tumor volume matched pairs (n=7 for each treatment group for each cell line). Drug or vehicle is administered by inter-peritoneal injection in 10% DMSO 90% sesame oil 2 to 3 times weekly for 5 weeks at 110 mg/kg to all mice harboring H358 xenografts or 3 times per week by gavage in 12.5% Cremophor EL, 12.5% DMSO as an aqueous suspension at 55 mg/kg to mice harboring A549 xenografts. Tumor volumes are monitored twice weekly by caliper measurements. Animals are weighed and observed during the five weeks of treatment. At the end point, mice are euthanized by CO<sub>2</sub> asphyxiation and cervical dislocation, and blood, tumors and major organs collected and weighed. Paired, unequal variance, one-tailed t-tests are performed across treatment groups using Excel software.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Cancer Cell. 2019 Apr 15;35(4):677-691.e10.
- ACS Nano. 2023 Jan 19.
- Proc Natl Acad Sci U S A. 2019 Feb 19;116(8):2961-2966.
- Acta Pharmacol Sin. 2021 Apr 13.
- Commun Biol. 2022 Sep 2;5(1):904.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

- 
- [1]. Wang L, et al. A small molecule modulates Jumonji histone demethylase activity and selectively inhibits cancer growth. Nat Commun, 2013. 4: p. 2035.
- [2]. Banelli B, et al. Small molecules targeting histone demethylase genes (KDMs) inhibit growth of temozolomide-resistant glioblastoma cells. Oncotarget. 2017 Apr 4.
- 

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA