



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



TPE-MI

Cat. No.:	HY-143218
CAS No.:	1245606-71-6
Molecular Formula:	C ₃₁ H ₂₃ NO ₂
Molecular Weight:	441.52
Target:	Huntingtin; Parasite
Pathway:	Neuronal Signaling; Anti-infection
Storage:	4°C, protect from light
	* In solvent : -80°C, 2 years; -20°C, 1 year (protect from light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 44.15 mg/mL (100.00 mM; Need ultrasonic)					
	Preparing Stock Solutions	<div><div>Solvent</div><div>Concentration</div></div>	Mass	1 mg	5 mg	10 mg
		1 mM		2.2649 mL	11.3245 mL	22.6490 mL
		5 mM		0.4530 mL	2.2649 mL	4.5298 mL
		10 mM		0.2265 mL	1.1325 mL	2.2649 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (5.66 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	TPE-MI (Tetraphenylethene maleimide) is a thiol probe for measuring unfolded protein load and proteostasis in cells. TPE-MI can report imbalances in proteostasis in induced pluripotent stem cell models of Huntington disease, as well as cells transfected with mutant Huntington exon 1 before the formation of visible aggregates. TPE-MI also detects protein damage following dihydroartemisinin research of the malaria parasites <i>Plasmodium falciparum</i> ^{[1][2]} .
IC ₅₀ & Target	Plasmodium
In Vitro	<p>TPE-MI is inherently non-fluorescent until it is conjugated to a thiol via the maleimide. TPE-MI fluorescence is activated upon labelling free cysteine thiols, normally buried in the core of globular proteins that are exposed upon unfolding^[1].</p> <p>TPE-MI (50 μM; 0-60 min) exhibits a homogeneous cytoplasmic labelling pattern in live HeLa cells, with a lower level of labelling in the nucleus and apparent concentration in the region of the ER, which was anticipated as a major location for protein synthesis and folding^[1].</p> <p>At the high expression level, the mutant 97Q form of Httex1 was associated with an elevated TPE-MI fluorescence signal relative to a non-disease-causing 25Q form of Httex1^[1].</p>

TPE-MI consists of the aggregation-induced emission (AIE) fluorogen tetraphenylethane (TPE) and the thiol-reactive group maleimide (MI), thereby possessing both AIE phenomenon and selective thiol reactivity^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Chen MZ, et al. A thiol probe for measuring unfolded protein load and proteostasis in cells. Nat Commun. 2017;8(1):474. Published 2017 Sep 7.
- [2]. Hu Q, et al. In Situ Monitored Vortex Fluidic-Mediated Protein Refolding/Unfolding Using an Aggregation-Induced Emission Bioprobe. Molecules. 2021;26(14):4273. Published 2021 Jul 14.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA