



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Olanzapine-d₈

Cat. No.:	HY-14541S2
CAS No.:	1093380-13-2
Molecular Formula:	C ₁₇ H ₁₂ D ₈ N ₄ S
Molecular Weight:	320.48
Target:	Apoptosis; Dopamine Receptor; Autophagy; mAChR; Mitophagy; 5-HT Receptor; Adrenergic Receptor; Isotope-Labeled Compounds
Pathway:	Apoptosis; GPCR/G Protein; Neuronal Signaling; Autophagy; Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Olanzapine-d ₈ is a deuterated labeled Olanzapine ^[1] . Olanzapine (LY170053) is a selective, orally active monoaminergic antagonist with high affinity binding to serotonin H ₁ , 5HT _{2A/2C} , 5HT ₃ , 5HT ₆ (K _i =7, 4, 11, 57, and 5 nM, respectively), dopamine D ₁₋₄ (K _i =11 to 31 nM), muscarinic M ₁₋₅ (K _i =1.9-25 nM), and adrenergic α ₁ receptor (K _i =19 nM). Olanzapine is an atypical antipsychotic ^{[2][3]} .
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs^[1]. Olanzapine binds weakly to GABAA, Benzodiazepine (BZD), and β-adrenergic receptors (K_i>10 μM) ^{[2][3]}. Olanzapine induces autophagy in human SH-SY5Y neuronal cell line^[4]. Olanzapine (1-100 μM for 144 h under serum starvation) results in a marked anti-proliferative effect in glioblastoma cell lines as well as glioma stem-like cells^[5]. Olanzapine also enhances Temozolomide (HY-17364)'s anti-tumor activity in glioblastoma cell lines^[5]. Olanzapine induces apoptosis and necrosis in glioblastoma cell lines^[5]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	Olanzapine (0.75, 1.5 and 3 mg/kg) evaluates body weight and periuterine fat mass, as well as insulin, non-esterified fatty acids, triglycerides, and glucose levels in mice^[6] MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Vucicevic L, et al. Autophagy inhibition uncovers the neurotoxic action of the antipsychotic drug olanzapine. *Autophagy*. 2014;10(12):2362-78.
- [2]. APPROVED AGREED-UPON LABELING.
- [3]. Karpel-Massler G, et al. Olanzapine inhibits proliferation, migration and anchorage-independent growth in human glioblastoma cell lines and enhances temozolomide's antiproliferative effect. *J Neurooncol*. 2015 Mar;122(1):21-33.
- [4]. Olanzapine for Injection, powder, for solution for intramuscular use.
- [5]. Coccarello R, et al. Chronic administration of olanzapine induces metabolic and food intake alterations: a mouse model of the atypical antipsychotic-associated adverse effects. *Psychopharmacology (Berl)*. 2006 Jul;186(4):561-71.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA