



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SGC-SMARCA-BRDVIII

Cat. No.:	HY-145446		
CAS No.:	1997319-84-2		
Molecular Formula:	C ₁₉ H ₂₅ N ₅ O ₃		
Molecular Weight:	371.43		
Target:	Epigenetic Reader Domain		
Pathway:	Epigenetics		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 25 mg/mL (67.31 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		2.6923 mL	13.4615 mL	26.9230 mL
		5 mM		0.5385 mL	2.6923 mL	5.3846 mL
		10 mM		0.2692 mL	1.3461 mL	2.6923 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (6.73 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (6.73 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (6.73 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	SGC-SMARCA-BRDVIII is a potent and selective inhibitor of SMARCA2/4 and PB1(5), with K _d s of 35 nM, 36 nM, and 13 nM, respectively. SGC-SMARCA-BRDVIII also inhibits PB1(2) and PB1(3), with K _d s of 3.7 and 2.0 μM, respectively. SGC-SMARCA-BRDVIII can block adipogenesis of 3T3-L1 murine fibroblasts ^{[1][2]} .
IC ₅₀ & Target	Kd: 35 nM (SMARCA2), 36 nM (SMARCA4), 13 nM (PB1(5)) ^[1]
In Vitro	SGC-SMARCA-BRDVIII (compound 22) (1 μM; 14 days) blocks adipocyte differentiation in 3T3-L1 murine fibroblasts ^[1] .

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Wanior M, et, al. Pan-SMARCA/PB1 Bromodomain Inhibitors and Their Role in Regulating Adipogenesis. J Med Chem. 2020 Dec 10;63(23):14680-14699.
- [2]. Mélin L, et, al. Design and Synthesis of LM146, a Potent Inhibitor of PB1 with an Improved Selectivity Profile over SMARCA2. ACS Omega. 2021 Aug 9;6(33):21327-21338.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA