



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CPPHA

Cat. No.:	HY-14612
CAS No.:	693288-97-0
Molecular Formula:	C ₂₂ H ₁₅ ClN ₂ O ₄
Molecular Weight:	406.82
Target:	mGluR
Pathway:	GPCR/G Protein; Neuronal Signaling
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (245.81 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		2.4581 mL	12.2904 mL	24.5809 mL
		5 mM		0.4916 mL	2.4581 mL	4.9162 mL
		10 mM		0.2458 mL	1.2290 mL	2.4581 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (6.15 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	CPPHA is potent and selective positive allosteric modulator (PAM) of the mGluR5 and mGluR1 (metabotropic glutamate receptor). CPPHA can potentiate responses of mGluR5 and mGluR1 to activation of these receptors. CPPHA is developed for the research of central nervous system disorders ^{[1][2]} .
IC ₅₀ & Target	mGlu5 Receptor
In Vitro	CPPHA alone has no agonist activity^[1]. CPPHA causes concentration-dependent potentiation of the response of human mGluR5 CHO cells to agonists^[1]. CPPHA potentiates threshold response to glutamate in fluorometric Ca²⁺ assays 7- to 8-fold with EC₅₀ values in the 400 to 800 nM range^[1]. CPPHA (20 μM; 15 minutes) potentiates the response to a subthreshold concentration of DHPG (HY-12598A) on ERK and cyclic-AMP responsive element-binding protein (CREB) activity, as well as NMDA receptor subunit NR1 phosphorylation in

cortical and hippocampal slices^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Western Blot Analysis^[3]

Cell Line:	Rat hippocampal cells
Concentration:	20 μ M
Incubation Time:	15 minutes
Result:	Potentiated the DHPG (20 μ M)-induced increase in phosphorylation of ERK and CREB, approaching levels seen with 100 μ M DHPG alone.

REFERENCES

- [1]. Liu F, et al. The effect of mGlu5 receptor positive allosteric modulators on signaling molecules in brain slices. *Eur J Pharmacol.* 2006 May 1;536(3):262-8.
- [2]. O'Brien JA, et al. A novel selective allosteric modulator potentiates the activity of native metabotropic glutamate receptor subtype 5 in rat forebrain. *J Pharmacol Exp Ther.* 2004 May;309(2):568-77.
- [3]. Zhang Y, et al. Allosteric potentiators of metabotropic glutamate receptor subtype 5 have differential effects on different signaling pathways in cortical astrocytes. *J Pharmacol Exp Ther.* 2005 Dec;315(3):1212-9.
- [4]. N-[4-Chloro-2-[(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)methyl]phenyl]-2-hydroxybenzamide (CPPHA) acts through a novel site as a positive allosteric modulator of group 1 metabotropic glutamate receptors. *Mol Pharmacol.* 2008 Mar;73(3):909-18.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA