



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

EBV lytic cycle inducer-1

Cat. No.:	HY-149577
CAS No.:	394668-43-0
Molecular Formula:	C ₁₄ H ₁₂ BrN ₃ O
Molecular Weight:	318.17
Target:	EBV
Pathway:	Anti-infection
Storage:	<div>Powder -20°C 3 years</div> <div> 4°C 2 years</div> <div>In solvent -80°C 6 months</div> <div> -20°C 1 month</div>



SOLVENT & SOLUBILITY

In Vitro	DMSO : 12.5 mg/mL (39.29 mM; ultrasonic and warming and heat to 60°C)					
	Preparing Stock Solutions	<div><div>Solvent</div><div>Concentration</div></div>	Mass	1 mg	5 mg	10 mg
		1 mM		3.1430 mL	15.7149 mL	31.4297 mL
		5 mM		0.6286 mL	3.1430 mL	6.2859 mL
		10 mM		0.3143 mL	1.5715 mL	3.1430 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 0.56 mg/mL (1.76 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 0.56 mg/mL (1.76 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 0.56 mg/mL (1.76 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	Epstein-Barr virus (EBV) lytic cycle inducer-1 Dp44mT (compound C7) is an iron-chelatoe-like compound. Dp44mT cooperates with HDAC inhibitor Romidespin (HY-15149) and SAHA to induce EBV lytic cycle. Dp44mT reactivates EBV lytic cycle by activating the ERK1/2-autophagy axis in epithelial cancers ^{[1][2]} .
In Vitro	<p>Dp44mT (compound C7) (0-80 μM; 48 h) induces lytic cycle in cell line-dependent manner, with higher toxicity in AGS-BX1 than in AGS^[1].</p> <p>Dp44mT (10 μM; 0-72 h) induces lytic cycle in a time-dependent manner^[1].</p>

Dp44mT (1.25-2.5 μ M; 24 h) cooperates with HDAC inhibitor Romidespin and SAHA to induce EBV lytic cycle^[1].
 Dp44mT (20 μ M; 48 h) leads to the EBV lytic cycle through induction of the ERK-autophagy axis^[2].
 MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Immunofluorescence^[1]

Cell Line:	AGS AGS-BX1
Concentration:	10 μ M
Incubation Time:	24 h, 48 h, 72 h
Result:	Resulted the expression of IE proteins Zta, Rta, and early EBV lytic protein BMRF1 peaking at 24h post treatment.

Immunofluorescence^[1]

Cell Line:	AGS-BX1
Concentration:	1.25 μ M, 2.5 μ M
Incubation Time:	24 h
Result:	Synergistically induced the expression of the viral IE protein Zta could together with 2.5 μ M of SAHA and 2.5 nM of Rmidepsin.

Immunofluorescence^[2]

Cell Line:	HA cells
Concentration:	20 μ M
Incubation Time:	24 h
Result:	A significantly lower expression of Zta was observed in cells treated with the iron-precomplexed C7 when compared to cells treated with C7 with 41% higher.

Cell Proliferation Assay^[1]

Cell Line:	AGS, AGS-BX1
Concentration:	0 μ M, 1.25 μ M, 2.5 μ M, 5 μ M, 10 μ M, 20 μ M, 40 μ M, 80 μ M
Incubation Time:	48 h
Result:	Displayed significantly higher toxicity to the EBV-positive cell line AGS-BX1 than the EBV-negative counterpart.

REFERENCES

[1]. Chung King Choi, et al. Identification of Novel Small Organic Compounds with Diverse Structures for the Induction of Epstein-Barr Virus (EBV) Lytic Cycle in EBV-Positive Epithelial Malignancies. PLoS One. 2015 Dec 30;10(12):e0145994.

[2]. Stephanie Pei Tung Yiu, et al. Intracellular Iron Chelation by a Novel Compound, C7, Reactivates Epstein-Barr Virus (EBV) Lytic Cycle via the ERK-Autophagy Axis in EBV-Positive Epithelial Cancers Cancers 2018 Dec; 10(12): 505.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA