



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DNQX

Cat. No.:	HY-15067
CAS No.:	2379-57-9
Molecular Formula:	C ₈ H ₄ N ₄ O ₆
Molecular Weight:	252.14
Target:	iGluR
Pathway:	Membrane Transporter/Ion Channel; Neuronal Signaling
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 35 mg/mL (138.81 mM)
H₂O : < 0.1 mg/mL (ultrasonic;warming;heat to 60°C) (insoluble)
* "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		3.9661 mL	19.8303 mL	39.6605 mL
	5 mM		0.7932 mL	3.9661 mL	7.9321 mL
	10 mM		0.3966 mL	1.9830 mL	3.9661 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.08 mg/mL (8.25 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: 2.08 mg/mL (8.25 mM); Suspended solution; Need ultrasonic

BIOLOGICAL ACTIVITY

Description

DNQX (FG 9041), a quinoxaline derivative, is a selective, potent competitive non-NMDA glutamate receptor antagonist (IC₅₀s = 0.5, 2 and 40 μM for AMPA, kainate and NMDA receptors, respectively)^[1].

IC₅₀ & Target

Kainate Receptor

In Vitro

DNQX selectively depolarizes thalamic reticular nucleus (TRN) neurons^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

DNQX, a specific AMPA receptor antagonist, given as either a 5 mg/kg or 10 mg/kg intraperitoneal dose or into the lateral cerebral ventricle (5 µl of 0.5 mg/ml) significantly diminishes phencyclidine (PCP) (40 mg/kg) and ketamine (80, 100, 120 mg/kg) hsp70 induction in the posterior cingulate and retrosplenial cortex^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cell Death Discov. 2020 Sep 17;6:87.
- Neural Regen Res. 2022 Jan;17(1):178-184.
- J Physiol. 2023 Jul 8.
- J Psychiatr Res. 2023 May 17;163:180-194.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Honoré T, et al. Quinoxalinediones: potent competitive non-NMDA glutamate receptor antagonists. Science. 1988;241(4866):701-703.

[2]. Lee SH, et al. Selective excitatory actions of DNQX and CNQX in rat thalamic neurons. J Neurophysiol. 2010;103(4):1728-1734.

[3]. Sharp JW, et al. DNQX inhibits phencyclidine (PCP) and ketamine induction of the hsp70 heat shock gene in the rat cingulate and retrosplenial cortex. Brain Res. 1995;687(1-2):114-124.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA