



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## SY-LB-35

|                    |                                                                                                    |
|--------------------|----------------------------------------------------------------------------------------------------|
| Cat. No.:          | HY-150795                                                                                          |
| CAS No.:           | 2603461-70-5                                                                                       |
| Molecular Formula: | C <sub>15</sub> H <sub>11</sub> N <sub>3</sub> O                                                   |
| Molecular Weight:  | 249.27                                                                                             |
| Target:            | TGF-beta/Smad; PI3K; Akt; ERK; JNK                                                                 |
| Pathway:           | Stem Cell/Wnt; TGF-beta/Smad; PI3K/Akt/mTOR; MAPK/ERK Pathway                                      |
| Storage:           | Powder    -20°C    3 years<br>4°C    2 years<br>In solvent   -80°C    6 months<br>-20°C    1 month |



## SOLVENT & SOLUBILITY

|                                                                               |                                                                                                                                           |                                                       |      |           |            |            |
|-------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------|------|-----------|------------|------------|
| In Vitro                                                                      | DMSO : 100 mg/mL (401.17 mM; Need ultrasonic)                                                                                             |                                                       |      |           |            |            |
|                                                                               | Preparing Stock Solutions                                                                                                                 | <div><div>Solvent</div><div>Concentration</div></div> | Mass | 1 mg      | 5 mg       | 10 mg      |
|                                                                               |                                                                                                                                           | 1 mM                                                  |      | 4.0117 mL | 20.0586 mL | 40.1171 mL |
|                                                                               |                                                                                                                                           | 5 mM                                                  |      | 0.8023 mL | 4.0117 mL  | 8.0234 mL  |
|                                                                               |                                                                                                                                           | 10 mM                                                 |      | 0.4012 mL | 2.0059 mL  | 4.0117 mL  |
| Please refer to the solubility information to select the appropriate solvent. |                                                                                                                                           |                                                       |      |           |            |            |
| In Vivo                                                                       | 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline<br>Solubility: ≥ 2.5 mg/mL (10.03 mM); Clear solution |                                                       |      |           |            |            |
|                                                                               | 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)<br>Solubility: ≥ 2.5 mg/mL (10.03 mM); Clear solution            |                                                       |      |           |            |            |

## BIOLOGICAL ACTIVITY

|                           |                                                                                                                                                                                                                                                                                                                                                                                                                 |
|---------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Description               | SY-LB-35 is a potent bone morphogenetic protein (BMP) receptor agonist. SY-LB-35 can stimulate significant increases in cell number and cell viability in the C2C12 myoblast cell line, and causes shifts towards the S and G2/M phases of the cell cycle. SY-LB-35 stimulates canonical Smad and non-canonical PI3K/Akt, ERK, p38 and JNK intracellular signaling pathways <sup>[1]</sup> .                    |
| IC <sub>50</sub> & Target | BMP <sup>[1]</sup>                                                                                                                                                                                                                                                                                                                                                                                              |
| In Vitro                  | SY-LB-35 (0.01-1000 μM; 24 h) stimulates significant increases in cell number and cell viability in the C2C12 myoblast cell line <sup>[1]</sup> .<br><br>SY-LB-35 (0.01-10 μM; 30 min or 15 min) stimulates Smad phosphorylation and nuclear translocation, activates the PI3K/Akt pathway and direct the cytoplasmic localization of p-Akt, stimulates the phosphorylation and activation of PI3K in the C2C12 |

cells<sup>[1]</sup>.

SY-LB-35 (0.01-10  $\mu$ M; 24 h) causes the cell cycle shifting to S and G2/M phases in the C2C12 cells<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Cell Viability Assay<sup>[1]</sup>

|                  |                                                                     |
|------------------|---------------------------------------------------------------------|
| Cell Line:       | C2C12 cells                                                         |
| Concentration:   | 0.01, 0.1, 1, 10, 100 and 1000 $\mu$ M                              |
| Incubation Time: | 24 h                                                                |
| Result:          | Stimulated significant increases in cell number and cell viability. |

#### Western Blot Analysis<sup>[1]</sup>

|                  |                                                                                                                                                                                                                       |
|------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Cell Line:       | C2C12 cells                                                                                                                                                                                                           |
| Concentration:   | 0.01, 0.1, 1 and 10 $\mu$ M                                                                                                                                                                                           |
| Incubation Time: | 30 or 15 min                                                                                                                                                                                                          |
| Result:          | Stimulated Smad phosphorylation and nuclear translocation, activated the PI3K/Akt pathway and direct the cytoplasmic localization of p-Akt, stimulated the phosphorylation and activation of PI3K in the C2C12 cells. |

#### Cell Cycle Analysis<sup>[1]</sup>

|                  |                                                      |
|------------------|------------------------------------------------------|
| Cell Line:       | C2C12 cells                                          |
| Concentration:   | 0.01, 0.1, 1 and 10 $\mu$ M                          |
| Incubation Time: | 24 h                                                 |
| Result:          | Caused the cell cycle shifting to S and G2/M phases. |

## REFERENCES

[1]. Najafi S, et al. Discovery of a novel class of benzimidazoles as highly effective agonists of bone morphogenetic protein (BMP) receptor signaling. Sci Rep. 2022 Jul 15;12(1):12146.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA