



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



RP-6685

Cat. No.:	HY-151462
CAS No.:	2832047-80-8
Molecular Formula:	C ₂₂ H ₁₄ F ₇ N ₅ O
Molecular Weight:	497.37
Target:	DNA/RNA Synthesis
Pathway:	Cell Cycle/DNA Damage
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (201.06 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM	2.0106 mL	10.0529 mL	20.1058 mL	
		5 mM	0.4021 mL	2.0106 mL	4.0212 mL	
		10 mM	0.2011 mL	1.0053 mL	2.0106 mL	
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% corn oil					
	Solubility: ≥ 2.08 mg/mL (4.18 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	RP-6685 is a potent, selective and orally active DNA polymerase theta (Polθ) inhibitor with an IC ₅₀ value of 5.8 nM (PicoGreen assay). RP-6685 shows antitumor efficacy in mouse tumor xenograft model ^[1] . RP-6685 is a click chemistry reagent, it contains an Alkyne group and can undergo copper-catalyzed azide-alkyne cycloaddition (CuAAC) with molecules containing Azide groups.
IC ₅₀ & Target	IC ₅₀ : 5.8 nM (Polθ) ^[1]
In Vitro	RP-6685 is extremely potent with an IC ₅₀ of 550 pM against the pol activity of full-length Polθ and inactive on the ATPase activity ^[1] . ?RP-6685 inhibits Polθ in HEK293 LIG4 ^{-/-} cellswith an IC ₅₀ of 0.94 μM ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

RP-6685 (80 mg/kg; p.o.; BID for 21 days) exhibits potent antitumor efficacy in BRCA2-deficient HCT116 mice^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Female CD1 nude mice (HCT116 BRCA2 ^{+/+} and BRCA2 ^{-/-} xenograft tumor models) ^[1]
Dosage:	80 mg/kg
Administration:	p.o.; BID for 21 days
Result:	Showed tumor regression during the first 8 days of treatment in BRCA2 ^{-/-} HCT116 model, while did not inhibit tumor growth in BRCA2 ^{+/+} HCT116 tumors mice.

Animal Model:	CD1 mice (20-30 g) ^[1]			
Dosage:	2.5 mg/kg			
Administration:	i.v. or p.o.; single dosage			
Result:				
	CL (mL/min/kg)	V _{dss} (L/kg)	t _{1/2} (h)	F (%)
	36.8	1.1	0.4	66

REFERENCES

[1]. Bubenik M, et al. Identification of RP-6685, an Orally Bioavailable Compound that Inhibits the DNA Polymerase Activity of Pol θ . J Med Chem. 2022 Sep 20.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA