



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

JY-2

Cat. No.:	HY-153347
CAS No.:	339103-05-8
Molecular Formula:	C ₁₃ H ₇ Cl ₂ N ₃ O
Molecular Weight:	292.12
Target:	FOXO
Pathway:	Metabolic Enzyme/Protease
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 100 mg/mL (342.33 mM; ultrasonic and warming and heat to 60°C)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		3.4233 mL	17.1163 mL	34.2325 mL
	5 mM		0.6847 mL	3.4233 mL	6.8465 mL
	10 mM		0.3423 mL	1.7116 mL	3.4233 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

JY-2 is a moderately selective and orally active Forkhead transcription factor forkhead box O1 (FoxO1) inhibitor that inhibits FoxO1 transcriptional activity with an IC₅₀ of 22 μM. JY-2 shows moderate inhibition against FoxO3a and FoxO4. JY-2 shows anti-diabetic activity^[1].

IC₅₀ & Target

22 μM (FoxO1 transcriptional activity)^[1]

In Vitro

JY-2 (10-100 μM; 24 h) reduces palmitic acid (PA; HY-N0830)-induced lipotoxicity in HepG2 and INS-1 cells^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Real Time qPCR^[1]

Cell Line:	HepG2 and INS-1 cells
Concentration:	10, 50 and 100 μM
Incubation Time:	24 h

Result:	Reduced palmitic acid (PA)-induced G6Pase and PEPCK mRNA expression. Inhibited PA-induced lipid accumulation. Reduced PA-induced mRNA expression of ER stress markers (ATF3, CHOP and GRP78).
Western Blot Analysis ^[1]	
Cell Line:	HepG2 cells
Concentration:	10, 50 and 100 μ M
Incubation Time:	4 h; in the presence of PA (500 μ M)
Result:	Increased p-FoxO1 levels in the whole cell lysate with a concurrent reduction in nuclear FoxO1 levels.

In Vivo

JY-2 (50-200 mg/kg; oral; 3 times for two days or daily for 4 weeks) shows anti-diabetic effects in mice^[1]. Pharmacokinetic parameters of JY-2^[1]

Parameters	i.v. (20 mg/kg)	p.o. (50 mg/kg)
AUC _{all} (ng·h/mL)	5017 $\pm$ 1038	12270 $\pm$ 2775
AUC _{inf.obs} (ng·h/mL)	5030 $\pm$ 1037	12400 $\pm$ 2753
C _{max} (ng/mL)	10790 $\pm$ 3269	6826 $\pm$ 2342
T _{max} (h)	0.1 $\pm$ 0.1	0.8 $\pm$ 0.7
T _{1/2} (h)	0.8 $\pm$ 0.2	1.3 $\pm$ 0.4
MRT _{inf.obs} (h)	0.7 $\pm$ 0.1	2.0 $\pm$ 0.1
F (%)		97.8

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	C57BL/6J mice ^[1]
Dosage:	50, 100, 200 mg/kg
Administration:	Oral, three times for two days (9:00 AM, 7:00 PM, 9:00 AM on the next day)
Result:	Improved glucose tolerance. Significantly reduced the expression of G6Pase and PEPCK mRNA in the liver. Enhanced mRNA expression of insulin and PDX-1 in the pancreas.
Animal Model:	db/db mice and C57BL/6J mice, high fat-diet-induced obese diabetic (DIO) model ^[1]
Dosage:	50, 100 mg/kg
Administration:	Oral, once daily for 4 weeks

Result:	Decreased the levels of fasting blood glucose, improved glucose tolerance. The expression of ColIV, a fibrosis marker, was also lowered.
---------	--

Animal Model:	C57BL/6J mice ^[1]
---------------	------------------------------

Dosage:	20 mg/kg or 50 mg/kg
---------	----------------------

Administration:	IV or PO (Pharmacokinetic Analysis)
-----------------	-------------------------------------

Result:	Showed an overall good pharmacokinetic profile.
---------	---

REFERENCES

[1]. Choi HE, et al. Novel FoxO1 inhibitor, JY-2, ameliorates palmitic acid-induced lipotoxicity and gluconeogenesis in a murine model. Eur J Pharmacol. 2021 May 15;899:174011.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA