



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Vesatolimod

Cat. No.:	HY-15601
CAS No.:	1228585-88-3
Molecular Formula:	C <sub>22</sub> H <sub>30</sub> N <sub>6</sub> O <sub>2</sub>
Molecular Weight:	410.51
Target:	Toll-like Receptor (TLR); Apoptosis; HBV; HCV; HIV
Pathway:	Immunology/Inflammation; Apoptosis; Anti-infection
Storage:	Powder    -20°C    3 years 4°C    2 years In solvent   -80°C    6 months -20°C    1 month



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : ≥ 16.67 mg/mL (40.61 mM)  
 \* "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.4360 mL	12.1800 mL	24.3599 mL
	5 mM		0.4872 mL	2.4360 mL	4.8720 mL
	10 mM		0.2436 mL	1.2180 mL	2.4360 mL

Please refer to the solubility information to select the appropriate solvent.

#### In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
Solubility: ≥ 1.67 mg/mL (4.07 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
Solubility: 1.67 mg/mL (4.07 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 90% corn oil  
Solubility: ≥ 1.67 mg/mL (4.07 mM); Clear solution

### BIOLOGICAL ACTIVITY

Description	Vesatolimod (GS-9620) is a potent, selective and orally active agonist of Toll-Like Receptor (TLR7) with an EC <sub>50</sub> of 291 nM.
IC <sub>50</sub> & Target	EC <sub>50</sub> : 291 nM (TLR7), 9 μM (TLR8) <sup>[3]</sup>
In Vitro	Vesatolimod (GS-9620) rapidly internalizes into cells and preferentially localizes to and signals from endo-lysosomal compartments. To test this hypothesis, the kinetics of cellular uptake of the compound in Daudi cells using tritiated

Vesatolimod ( $^3\text{H}$ -GS-9620) is measured. The kinetics of  $^3\text{H}$ -GS-9620 accumulation is rapid, reaching concentration-dependent steady-state equilibrium in approximately thirty minutes. Measured intracellular concentration of  $^3\text{H}$ -Vesatolimod is 5-fold higher than the extracellular concentration of  $^3\text{H}$ -GS-9620 used to treat cells. Increases in intracellular  $^3\text{H}$ -Vesatolimod concentrations are roughly proportional with increasing concentrations of  $^3\text{H}$ -GS-9620<sup>[1]</sup>. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### In Vivo

Single oral doses of Vesatolimod (GS-9620) at 0.3 and 1 mg/kg in uninfected chimpanzees demonstrates a dose- and exposure-related induction of serum IFN- $\alpha$ , select cytokines/chemokines, and IFN-stimulated genes (ISG) in the peripheral blood and liver. Following oral administration at 0.3 (n=3), and 1 mg/kg (n=3 and n=4), Vesatolimod (GS-9620)  $C_{\text{max}}$  is  $3.6 \pm 3.5$ ,  $36.8 \pm 34.5$ , and  $55.4 \pm 81.0$  nM, respectively. Peak serum IFN responses occur at 8 h post-dose. The mean peak levels of induced serum IFN- $\alpha$  are 66 and 479 pg/mL at doses of 0.3 and 1 mg/kg, respectively. Vesatolimod (GS-9620) treatment induces ISG transcripts including ISG15, OAS-1, MX1, IP-10 (CXCL10), and I-TAC (CXCL11) in peripheral blood mononuclear cells (PBMC) at 0.3 mg/kg and in both PBMC and the liver at 1 mg/kg<sup>[2]</sup>. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

#### Cell Assay<sup>[1]</sup>

Daudi cells are incubated for indicated times with varying concentrations [ $^3\text{H}$ ]Vesatolimod (GS-9620) (0.7  $\mu\text{Ci/mL}$ ). Cell associated radioactivity is extracted with ice cold 80% ethanol and measured using liquid scintillation counting. The total amount of Vesatolimod in cells is calculated from a calibration curve for Vesatolimod (GS-9620) mass versus radioactivity. Cell volume is measured<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Animal Administration<sup>[2]</sup>

Chimpanzee<sup>[2]</sup>  
Chimpanzees are used. The trial design includes 4 weeks of pre-study evaluation (Day -28, -13 and just prior to first dose) and two cycles of oral Vesatolimod (GS-9620) treatment every other day three times per week for 4 weeks with one cycle at 1 mg/kg, and, after a one week rest, a second cycle at 2 mg/kg. Animals are also intensely monitored for 14 weeks after treatment to assess tolerability and durability of response.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Nat Biomed Eng. 2018 Aug;2(8):578-588.
- Antiviral Res. 2022 Jul 19;205:105384.
- Antiviral Res. 2018 May;153:39-48.
- Vaccine. 2018 Feb 1;36(6):794-801.
- ACS Infect Dis. 2022 Dec 6.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Rebbapragada I, et al. Molecular Determinants of GS-9620-Dependent TLR7 Activation. PLoS One. 2016 Jan 19;11(1):e0146835.

[2]. Lanford RE, et al. GS-9620, an Oral Agonist of Toll-Like Receptor-7, Induces Prolonged Suppression of Hepatitis B Virus in Chronically Infected Chimpanzees. Gastroenterology. 2013 Feb 13. pii: S0016-5085(13)00169-8.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA