



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Ellipticine hydrochloride

Cat. No.:	HY-15753A
CAS No.:	5081-48-1
Molecular Formula:	C ₁₇ H ₁₅ ClN ₂
Molecular Weight:	282.77
Target:	Topoisomerase
Pathway:	Cell Cycle/DNA Damage
Storage:	4°C, sealed storage, away from moisture
	* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 5.8 mg/mL (20.51 mM; Need ultrasonic and warming)					
	H ₂ O : < 0.1 mg/mL (ultrasonic;warming;heat to 60°C) (insoluble)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		3.5364 mL	17.6822 mL	35.3644 mL
		5 mM		0.7073 mL	3.5364 mL	7.0729 mL
		10 mM		0.3536 mL	1.7682 mL	3.5364 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 0.84 mg/mL (2.97 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 0.84 mg/mL (2.97 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil					
	Solubility: ≥ 0.84 mg/mL (2.97 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	Ellipticine (NSC 71795) hydrochloride is a potent antineoplastic agent; inhibits DNA topoisomerase II activities.
IC ₅₀ & Target	Topoisomerase II
In Vitro	Ellipticine (NSC 71795) is a potent antineoplastic agent exhibiting the multimodal mechanism of its action. The mechanisms of Ellipticine (NSC 71795) antitumor, mutagenic and cytotoxic activities are suggested to be intercalation into DNA and inhibition of DNA topoisomerase II activity. Another mode of Ellipticine (NSC 71795) action is the formation of covalent DNA adducts mediated by its oxidation with cytochromes P450 (CYP) and peroxidases ^[1] . Ellipticine (NSC 71795) can also act as

an inhibitor or inducer of biotransformation enzymes, thereby modulating its own metabolism leading to its genotoxic and pharmacological effects. Treatment of cells with Ellipticine (NSC 71795) results in inhibition of cell growth and proliferation. This effect is associated with formation of two covalent Ellipticine (NSC 71795)-derived DNA adducts^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Ellipticine (NSC 71795) treatment results in ellipticine-derived DNA adduct generation in several healthy organs (liver, kidney, lung, spleen, breast, heart and brain) and in DNA of mammary adenocarcinoma. The levels of Ellipticine (NSC 71795)-derived DNA adducts generated in these adenocarcinomas are almost 2-fold higher than in normal healthy mammary tissue. The induced expression of cytochrome b₅ protein in liver of rats treated with Ellipticine (NSC 71795) suggests that cytochrome b₅ may modulate the CYP-mediated bioactivation and detoxification of Ellipticine (NSC 71795)^[3]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[2]

The cytotoxicity of Ellipticine (NSC 71795) is determined by MTT test. Ellipticine (NSC 71795) is dissolved in DMSO (1 mM) and diluted in culture medium to final concentrations of 0, 0.1, 1, 5 or 10 µM. Cells in exponential growth are seeded at 1×10⁴ per well in a 96-well microplate. After incubation the MTT solution is added, the microplates are incubated for 4 hours and cells lysed in 50% N,N-dimethylformamide containing 20% of sodium dodecyl sulfate (SDS), pH 4.5. The absorbance at 570 nm is measured. The mean absorbance of medium controls is subtracted as a background. The viability of control cells is taken as 100% and the values of treated cells are calculated as a percentage of control. The IC₅₀ values are calculated using linear regression of the dose-log response curves^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- J Adv Res. 2023 Jun 6;S2090-1232(23)00148-0.
- J Cell Biol. 2023 Jan 2;222(1):e202202110.
- Front Cell Dev Biol. 2019 Sep 20;7:204.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Stiborova M, et al. Molecular mechanisms of antineoplastic action of an anticancer drug ellipticine. Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub. 2006 Jul;150(1):13-23.

[2]. Stiborova M, et al. Ellipticine cytotoxicity to cancer cell lines - a comparative study. Interdiscip Toxicol. 2011 Jun;4(2):98-105.

[3]. Stiborova M, et al. The anticancer drug ellipticine activated with cytochrome P450 mediates DNA damage determining its pharmacological efficiencies: studies with rats, Hepatic Cytochrome P450 Reductase Null (HRN?) mice and pure enzymes. Int J Mol Sci. 2014 Dec 25;16(1):284-306.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA