



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TRITC-dextran, MW 40000

Cat. No.:	HY-158082B
Molecular Weight:	40000
Target:	Fluorescent Dye
Pathway:	Others
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)

TRITC-dextran, MW 40000

SOLVENT & SOLUBILITY

In Vitro

H₂O : 50 mg/mL (1.25 mM; Need ultrasonic)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		0.0250 mL	0.1250 mL	0.2500 mL
	5 mM		---	---	---
	10 mM		---	---	---

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

TRITC-dextran, MW 40000 (Tetramethyl rhodamine isothiocyanate glucan, MW 40000) is a fluorescent dye, with the molecular weight of 40 kD. TRITC-dextran MW 40000 exhibits an excitation wavelength of 555 nm. TRITC-dextran MW 40000 is vessel penetrate, which could label blood plasma to visualize the vasculature. TRITC-dextran MW 40000 is utilized in drug delivery for the stability of TRITC over a wide pH range (i.e. pH 2–11) and resistance to photo-bleaching^{[1][2][3]}.

REFERENCES

- [1]. Tran NBNN, et al., Gradient-dependent release of the model drug TRITC-dextran from FITC-labeled BSA hydrogel nanocarriers in the hair follicles of porcine ear skin. *Eur J Pharm Biopharm.* 2017 Jul;116:12-16.
- [2]. Kawoos U, et al., N-acetylcysteine Ameliorates Blast-Induced Changes in Blood-Brain Barrier Integrity in Rats. *Front Neurol.* 2019 Jun 26;10:650.
- [3]. Somaratne G, et al., In-situ disintegration of egg white gels by pepsin and kinetics of nutrient release followed by time-lapse confocal microscopy[J]. *Food hydrocolloids*, 2020, 98: 105228.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA