



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

LDN-212854

Cat. No.:	HY-15897		
CAS No.:	1432597-26-6		
Molecular Formula:	C ₂₅ H ₂₂ N ₆		
Molecular Weight:	406		
Target:	TGF-β Receptor		
Pathway:	TGF-beta/Smad		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 30 mg/mL (73.89 mM)
 * "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.4631 mL	12.3153 mL	24.6305 mL
	5 mM		0.4926 mL	2.4631 mL	4.9261 mL
	10 mM		0.2463 mL	1.2315 mL	2.4631 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
 Solubility: 2.5 mg/mL (6.16 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
 Solubility: 2.5 mg/mL (6.16 mM); Suspended solution; Need ultrasonic

BIOLOGICAL ACTIVITY

Description

LDN-212854 is a bone morphogenetic protein (BMP) inhibitor that potently inhibits ALK2 (IC₅₀: 1.3 nM). LDN-212854 also inhibits ALK1 (IC₅₀: 2.40 nM). LDN-212854 can be used in the research of fibrodysplasia ossificans progressive and cancers, such as hepatocellular carcinoma (HCC)^{[1][2]}.

IC₅₀ & Target

ACVR1 1.3 nM (IC ₅₀)	ALK1 2.4 nM (IC ₅₀)	BMPRI1A 85.8 nM (IC ₅₀)	ALK4 2133 nM (IC ₅₀)
ALK5 9276 nM (IC ₅₀)			

In Vitro

LDN-212854 (0-3.815 μ M) blocks the phosphorylation of SMAD1/5/8 induced by BMP7 in BMPR2^{-/-} cells^[1].
LDN-212854 (2.5 μ M, 5 days) inhibits cell proliferation in Huh7 and MT cells^[2].
LDN-212854 (0.5 μ M, 48 h) suppresses ID1 and EpCAM expression in Huh7 and MT cells^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.
Western Blot Analysis^[1]

Cell Line:	BMPR2-deficient pulmonary vascular smooth muscle cells
Concentration:	0, 1, 3, 6, 16, 39, 98, 244, 610, 1530, 3815 nM
Incubation Time:	
Result:	Inhibited the phosphorylation of SMAD1/5/8 induced by BMP7 with an IC ₅₀ value of 37 nM.

In Vivo

LDN-212854 (intraperitoneal injection, 6 mg/kg, twice daily for 4 weeks) potently inhibits heterotopic ossification in an inducible transgenic mutant ALK2 mouse model of fibrodysplasia ossificans progressiva^[1].
LDN-212854 (intraperitoneal injection, 6 mg/kg, twice daily for 10-14 days) suppresses HCC tumor progression through repression of ID1 in HCC xenografts model^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Murine inducible transgenic ALK2Q207D model of heterotopic ossification ^[1]
Dosage:	6 mg/kg
Administration:	Intraperitoneal injection , twice daily for 4 weeks
Result:	Prevented the formation of heterotopic bone and preserved limb range of motion with minimal or no impairment in the majority of mice.

Animal Model:	HCC xenografts (Huh7 or MT cell) ^[1]
Dosage:	6 mg/kg
Administration:	Intraperitoneal injection, twice daily for 10-14 days.
Result:	Inhibited tumor growth and showed less spheroid/colony formation ability than PBS-treated tumor cells.

CUSTOMER VALIDATION

- Adv Sci (Weinh). 2024 Jan 16:e2306499.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Han Chen, et al. BMP9-ID1 signaling promotes EpCAM-positive cancer stem cell properties in hepatocellular carcinoma. Mol Oncol. 2021 Aug;15(8):2203-2218.
- [2]. Mohedas AH, et al. Development of an ALK2-biased BMP type I receptor kinase inhibitor. ACS Chem Biol. 2013;8(6):1291-302.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA