



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ML-T7

Cat. No.:	HY-163028		
CAS No.:	459789-75-4		
Molecular Formula:	C ₂₇ H ₁₇ Cl ₂ NO ₅		
Molecular Weight:	506.33		
Target:	Tim3		
Pathway:	Immunology/Inflammation		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 66.67 mg/mL (131.67 mM; Need ultrasonic)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		1.9750 mL	9.8750 mL	19.7500 mL
	5 mM		0.3950 mL	1.9750 mL	3.9500 mL
	10 mM		0.1975 mL	0.9875 mL	1.9750 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

ML-T7 is a potent Tim-3 inhibitor. ML-T7 blocks Tim-3 interactions with PtdSer and CEACAM1. ML-T7 not only enhances the antitumor activity of adoptive transfer therapy with cytotoxic T lymphocytes (CTLs) and CAR T cells but also increases the effector function of T cell. ML-T7 promotes NK cells' killing activity against tumor cells and DC antigen-presenting capacity. ML-T7 directly exerts antitumor efficacy in preclinical tumor models either alone or in combination with Nivolumab (HY-P9903A). ML-T7 can be used for tumor immunotherapy research^[1].

In Vitro

ML-T7(10 μM, 0-6 days) enhances TCR/STAT5 signaling and promotes CD8⁺ cell antitumor activity through Tim-3^[1].
 ML-T7(10 μM, 24 h) promotes DC maturation and function through Tim-3 and Tim-4, enhances DC antigen-presenting ability^[1].
 ML-T7(10 μM, 24 h) enhances CTL activation and cytokine production, decreases apoptosis of CTLs^[1].
 ML-T7(10 μM, 48 h) significantly enhances the production of IFN-γ, TNF-α, CD107a, and granzyme B in human NK cell line NK92 cells^[1].
 MCE has not independently confirmed the accuracy of these methods. They are for reference only.
 Western Blot Analysis^[1]

Cell Line:	CD8 ⁺ cell
Concentration:	10 μ M
Incubation Time:	0-6 days
Result:	Increased the phosphorylation of phospholipase C- γ 1(PLC- γ 1), ZAP70, LCK, ERK1/2, and STAT5 upon anti-CD3/CD28 stimulation.

Cell Invasion Assay^[1]

Cell Line:	Bone Marrow Cells
Concentration:	10 μ M
Incubation Time:	24 h
Result:	Increased the expression of DC maturation markers.

In Vivo

ML-T7 (10-50 mg/kg; Intraperitoneal injection; every 2 days, 10 times) inhibits the growth of tumor and prolongs survival of HCC mice, without adverse effects on mice body weight^[1].

ML-T7 (20 mg/kg; Intraperitoneal injection; every 2 days, 10 times) synergizes with Nivolumab (HY-P9903A) to improve the antitumor efficacy of Nivolumab (HY-P9903A) antibodies, and effectively improves HCC mice survival^[1].

ML-T7 (50 mg/kg, Intraperitoneal injection; every 2 days for 3 weeks) shows a good safety profile in mice^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	10-week-old mice with spontaneous orthotopic HCC ^[1]
Dosage:	10-50 mg/kg
Administration:	Intraperitoneal injection (i.p.), every 2 days, 10 times
Result:	Increased CD8 ⁺ T cells in both tumor and spleen. Inhibited T cell exhaustion. Promoted the function of CTLs, NK cells, and DCs.

Animal Model:	10-week-old mice with orthotopic Akt/c-Myc HCC ^[1]
Dosage:	20 mg/kg
Administration:	Intraperitoneal injection (i.p.), every 2 days, 10 times
Result:	Indicated stronger antitumor activity when treated with ML-T7 and Nivolumab (HY-P9903A). Rejuvenated NK cells by the combination therapy. Inhibited the accumulation of MDSCs and Tregs.

REFERENCES

[1]. Ma S, et al. Identification of a small-molecule Tim-3 inhibitor to potentiate T cell-mediated antitumor immunotherapy in preclinical mouse models. Sci Transl Med. 2023 Nov 15;15(722):eadg6752.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA