



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SMIFH2

Cat. No.:	HY-16931
CAS No.:	340316-62-3
Molecular Formula:	C ₁₅ H ₉ BrN ₂ O ₃ S
Molecular Weight:	377.21
Target:	Arp2/3 Complex
Pathway:	Cytoskeleton
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 125 mg/mL (331.38 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM	2.6510 mL	13.2552 mL	26.5104 mL	
		5 mM	0.5302 mL	2.6510 mL	5.3021 mL	
	10 mM	0.2651 mL	1.3255 mL	2.6510 mL		
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)					
	Solubility: 2.08 mg/mL (5.51 mM); Suspended solution; Need ultrasonic					

BIOLOGICAL ACTIVITY

Description	SMIFH2 is a formin specific inhibitor. SMIFH2 inhibits actin polymerization by Formins and affects the actin cytoskeleton ^[1] .
IC ₅₀ & Target	Formin ^[1]
In Vitro	SMIFH2 (25 μM; 1-16 hours) induces dynamic cytoskeletal remodelling in U2OS cells ^[1] . SMIFH2 (25 μM) reduces p300, mDia2 and p53 levels in a proteasome-independent manner ^[1] . SMIFH2 reduces expression and activity of p53 through a post-transcriptional, proteasome-independent mechanism that influences remodelling of the cytoskeleton ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only. Western Blot Analysis ^[1]

Cell Line:	293T, A375, U2OS, and MDA-MB-231 cells
Concentration:	25 μ M
Incubation Time:	5 hours for 293T, U2OS, MDA-MB-231 cells; 2.5 hours for A375 cells
Result:	Downregulated mDia2, p53 and p300 protein levels.

CUSTOMER VALIDATION

- Theriogenology. 2023 Nov, 211, 40-48.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Tadamoto Isogai, et al. SMIFH2 has effects on Formins and p53 that perturb the cell cytoskeleton. Sci Rep. 2015 Apr 30;5:9802.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA