



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Alminoprofen

Cat. No.:	HY-17485
CAS No.:	39718-89-3
Molecular Formula:	C ₁₃ H ₁₇ NO ₂
Molecular Weight:	219.28
Target:	Phospholipase; COX
Pathway:	Metabolic Enzyme/Protease; Immunology/Inflammation
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (456.04 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		4.5604 mL	22.8019 mL	45.6038 mL
		5 mM		0.9121 mL	4.5604 mL	9.1208 mL
		10 mM		0.4560 mL	2.2802 mL	4.5604 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (11.40 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (11.40 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil					
	Solubility: ≥ 2.5 mg/mL (11.40 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	Alminoprofen (EB-382) is a nonsteroidal anti-inflammatory agent (NSAID) of the phenylpropionic acid class. Alminoprofen possesses a dual anti-inflammatory action, by inhibiting both secretory phospholipase A ₂ (sPLA ₂) and COX-2 ^[1] .	
IC ₅₀ & Target	sPLA ₂	COX-2
In Vitro	Alminoprofen possesses a dual anti-inflammatory action, by inhibiting both sPLA ₂ but also COX-2. Alminoprofen (from 0 to 50 μM) induces a dose-related inhibitory response on the levels of PGE ₂ formed by the CRL-1517 cell line ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

In Vivo

Alminoprofen (100-300 mg/kg; p.o.) exerts an anti-inflammatory activity^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Groups of Wistar Rats (Oedema of the paw) ^[1]
Dosage:	100 mg/kg
Administration:	P.o.
Result:	Exerted a statistically significant dose-dependent antagonist action lasting throughout the 2-hr observation period.

REFERENCES

[1]. Raguene-Nicol C, et al. Anti-inflammatory mechanism of alminoprofen: action on the phospholipid metabolism pathway. Biochem Pharmacol. 1999;57(4):433-443.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA