



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



EGFR-IN-12

Cat. No.:	HY-17499		
CAS No.:	879127-07-8		
Molecular Formula:	C ₂₁ H ₁₈ F ₃ N ₅ O		
Molecular Weight:	413.4		
Target:	EGFR; Apoptosis		
Pathway:	JAK/STAT Signaling; Protein Tyrosine Kinase/RTK; Apoptosis		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : 250 mg/mL (604.74 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass		1 mg	5 mg	10 mg
	Concentration				
	1 mM		2.4190 mL	12.0948 mL	24.1896 mL
	5 mM		0.4838 mL	2.4190 mL	4.8379 mL
	10 mM		0.2419 mL	1.2095 mL	2.4190 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

EGFR-IN-12 is a 4,6-disubstituted pyrimidine and is a potent, ATP-competitive, irreversible and highly selective EGFR inhibitor with an IC₅₀ of 21 nM. EGFR-IN-12 also inhibits mutant EGFR^{L858R} and EGFR^{L861Q} with IC₅₀s of 63 nM and 4 nM, respectively. EGFR-IN-12 displays strong selectivity for EGFR over HER4 (IC₅₀ = 7640 nM) and a panel of 55 other kinases. EGFR-IN-12 induces cells apoptosis and has antitumor activity^{[1][2]}.

IC₅₀ & Target

EGFR (WT) 21 nM (IC ₅₀)	EGFR ^{L858R} 63 nM (IC ₅₀)	EGFR ^{L861Q} 4 nM (IC ₅₀)	HER4 7640 nM (IC ₅₀)
--	--	---	-------------------------------------

In Vitro

EGFR-IN-12 (EGFR inhibitor 324674; 0-2 μM; 48 hours; HT29 and SW480 cells) treatment efficiently induces apoptosis at lower concentrations^[2].
 EGFR-IN-12 (EGFR inhibitor 324674; 0-3 μM; 3 hours; HT29 and SW480 cells) treatment inhibits EGFR activation and the downstream AKT signaling pathway in a dose-dependent manner^[2].
 EGFR-IN-12 (EGFR inhibitor 324674) inhibits HT29 and SW480 cell proliferation with with IC₅₀s of 1.96 μM and 1.04 μM, respectively^[2].
 Pretreatment of cells with EGFR-IN-12 (compound 1; 10 μM) results in complete inhibition of wild-type receptor

autophosphorylation in U-2OS cells. And the T766M mutant receptor is completely resistant to inhibition by EGFR-IN-12^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Apoptosis Analysis^[1]

Cell Line:	HT29 and SW480 cells
Concentration:	0 μ M, 1 μ M, 2 μ M
Incubation Time:	48 hours
Result:	Induced apoptosis in HT29 cells and SW480 cells.

Western Blot Analysis^[1]

Cell Line:	HT29 and SW480 cells
Concentration:	0 μ M, 0.1 μ M, 0.3 μ M, 1 μ M, 3 μ M
Incubation Time:	3 hours
Result:	Inhibited EGFR activation and the downstream AKT signaling pathway in a dose-dependent manner.

REFERENCES

[1]. Qiong Zhang, et al. Discovery of EGFR Selective 4,6-disubstituted Pyrimidines From a Combinatorial Kinase-Directed Heterocycle Library. J Am Chem Soc. 2006 Feb 22;128(7):2182-3.

[2]. Zhiwei Yu, et al. Novel Irreversible EGFR Tyrosine Kinase Inhibitor 324674 Sensitizes Human Colon Carcinoma HT29 and SW480 Cells to Apoptosis by Blocking the EGFR Pathway. Biochem Biophys Res Commun. 2011 Aug 12;411(4):751-6.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA