



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

OSU-T315

Cat. No.:	HY-18676
CAS No.:	2070015-22-2
Molecular Formula:	C ₃₀ H ₃₀ F ₃ N ₅ O
Molecular Weight:	533.59
Target:	Integrin; Autophagy; Apoptosis
Pathway:	Cytoskeleton; Autophagy; Apoptosis
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 260 mg/mL (487.27 mM)

* "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		1.8741 mL	9.3705 mL	18.7410 mL
	5 mM		0.3748 mL	1.8741 mL	3.7482 mL
	10 mM		0.1874 mL	0.9370 mL	1.8741 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.17 mg/mL (4.07 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.17 mg/mL (4.07 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

OSU-T315 (ILK-IN-1) is a small Integrin-linked kinase (ILK) inhibitor with an IC₅₀ of 0.6 μM, inhibiting PI3K/AKT signaling by dephosphorylation of AKT-Ser473 and other ILK targets (GSK-3β and myosin light chain)^[1]. OSU-T315 abrogates AKT activation by impeding AKT localization in lipid rafts and triggers caspase-dependent apoptosis in an ILK-independent manner^[2]. OSU-T315 causes cell death through apoptosis and autophagy^[1].

IC₅₀ & Target

IC₅₀: 0.6μM; Integrin-linked kinase (ILK) inhibitor^[1]

In Vitro

OSU-T315 (Compound 22; 0-5 μM; 24 hours) exhibits high in vitro potency against a panel of prostate and breast cancer cell lines with a IC₅₀ range of 1-2.5 μM^[1].

?OSU-T315 (0-2.5 μ M; 24 hours) can reduce YB-1, HER2, and EGFR expression; shows a modest suppressive effect on phosphorylated S6 levels, exhibits dose-dependent suppressive effects on the levels of phospho-ERK1/2 and phospho-p38, while that of phospho-JNK remains unaltered in PC-3 cell^[1].

?OSU-T315 (0-4 μ M; 24 hours) causes autophagy through ILK inhibition^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Western Blot Analysis^[1]

Cell Line:	PC-3 cells; MDA-MB-231 cells
Concentration:	1 μ M, 2 μ M, 3 μ M, 4 μ M; 0.5 μ M, 1 μ M, 1.5 μ M, 2 μ M, 2.5 μ M
Incubation Time:	24 hours
Result:	Exhibited a dose-dependent decreasing effect on the phosphorylation of pS6, ERKs, and p38 in PC-3 cells and MDA-MB-231 cells.

Cell Viability Assay^[1]

Cell Line:	Prostate cancer cells: LNCaP, PC-3; breast cancer cells: MDA-MB-231, MDA-MB-468, SKBR3, MCF-7; PrEC and MEC cells
Concentration:	0-5 μ M
Incubation Time:	24 hours
Result:	Suppressed cancer cells viability in breast and prostate cancer cells (IC (50), 1-2.5 μ M).

Apoptosis Analysis^[1]

Cell Line:	PC-3 cells
Concentration:	1 μ M, 2 μ M, 3 μ M, 4 μ M
Incubation Time:	24 hours
Result:	Induced accumulation of LC3-II and PARP cleavage.

In Vivo

OSU-T315 (Oral gavage; 25 mg/kg, 50 mg/kg; single daily; 35 days) has a suppressive effect of on PC-3 xenograft tumor growth^[1].

?No other obvious toxicity is observed in mice^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Male NCr athymic nude mice with PC-3 tumor xenografts
Dosage:	25 mg/kg; 50 mg/kg
Administration:	Oral gavage; single daily; 35 days
Result:	Resulted in suppression of tumor growth relative to the vehicle control after 35 days of treatment (48% and 62% suppression for 25 and 50 mg/kg, respectively).

CUSTOMER VALIDATION

- Theranostics. 2022 Jan 1;12(3):1173-1186.

-
- Br J Cancer. 2020 Aug;123(4):542-555.
 - PLoS Pathog. 2023 Mar 17;19(3):e1011241.
 - Biochim Biophys Acta Mol Basis Dis. 2020 Mar 1;1866(3):165625.
 - Colloids Surf B Biointerfaces. 27 November 2021, 112229.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Su-Lin Lee, et al Identification and Characterization of a Novel Integrin-Linked Kinase Inhibitor. J Med Chem. 2011 Sep 22; 54(18): 6364–6374
- [2]. Liu TM, et al. OSU-T315: a novel targeted therapeutic that antagonizes AKT membrane localization and activation of chronic lymphocytic leukemia cells. Blood. 2015 Jan 8;125(2):284-95.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA