



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

LLY-507

Cat. No.:	HY-19313
CAS No.:	1793053-37-8
Molecular Formula:	C ₃₆ H ₄₂ N ₆ O
Molecular Weight:	574.76
Target:	Histone Methyltransferase
Pathway:	Epigenetics
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 15 mg/mL (26.10 mM; Need ultrasonic and warming)				
	Preparing Stock Solutions	Solvent Concentration	Mass		
			1 mg	5 mg	10 mg
		1 mM	1.7399 mL	8.6993 mL	17.3986 mL
		5 mM	0.3480 mL	1.7399 mL	3.4797 mL
		10 mM	0.1740 mL	0.8699 mL	1.7399 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: 2.5 mg/mL (4.35 mM); Suspended solution; Need ultrasonic				
	2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (4.35 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	LLY-507 is a potent and selective inhibitor of protein-lysine methyltransferase SMYD2. LLY-507 potently inhibits the ability of SMYD2 to methylate p53 peptide with an IC ₅₀ <15 nM. LLY-507 serves as a valuable chemical probe to aid in the dissection of SMYD2 function in cancer and other biological processes ^[1] .
IC ₅₀ & Target	SMYD2 <15 nM (IC ₅₀)
In Vitro	LLY-507 is >100-fold selective for SMYD2 over 21 other methyltransferases including SMYD3 ^[1] . LLY-507 binds to the substrate channel of SMYD2 and supports the selectivity of its inhibition ^[1] . LY-507 is able to potently inhibit the methylation of H4 peptide by SMYD2 enzyme with an IC ₅₀ of 31 nM ^[1] .

LLY-507 (0.03-20 μ M; 28 hours) inhibits SMYD2-mediated methylation of p53 Lys³⁷⁰ in U2OS cells, with an IC₅₀ of 0.6 μ M^[1].
LLY-507 (0-20 μ M; 3-7 days) inhibits the proliferation of several ESCC, HCC, and breast cancer cell lines^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Proliferation Assay^[1]

Cell Line:	ESCC, HCC, breast cancer cell lines
Concentration:	0-20 μ M
Incubation Time:	3 days, 7 days
Result:	Inhibited tumor cell proliferation.

Western Blot Analysis^[1]

Cell Line:	HEK293 cells
Concentration:	0.03 μ M, 0.07 μ M, 0.15 μ M, 0.3 μ M, 0.6 μ M, 1.25 μ M, 2.5 μ M
Incubation Time:	28 hours
Result:	Inhibited SMYD2-mediated methylation of p53 Lys ³⁷⁰ in cells.

CUSTOMER VALIDATION

- Exp Mol Med. 2023 May 1.
- Cell Death Dis. 2022 Oct 21;13(10):890.
- Cell Death Dis. 2018 Jan 26;9(2):129.
- Cell Death Discov. 2022 Jun 6;8(1):274.
- Cells. 2022 Apr 8;11(8):1262.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Nguyen H, et al. LLY-507, a Cell-active, Potent, and Selective Inhibitor of Protein-lysine Methyltransferase SMYD2. J Biol Chem. 2015 May 29;290(22):13641-13653.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA