



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



MS023 dihydrochloride

Cat. No.: HY-19615B
CAS No.: 1992047-64-9
Molecular Formula: C₁₇H₂₇Cl₂N₃O
Molecular Weight: 360.32
Target: Histone Methyltransferase
Pathway: Epigenetics
Storage: 4°C, stored under nitrogen
 * In solvent : -80°C, 6 months; -20°C, 1 month (stored under nitrogen)



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 150 mg/mL (416.30 mM)
 H₂O : 33.33 mg/mL (92.50 mM; Need ultrasonic)
 * "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.7753 mL	13.8766 mL	27.7531 mL
	5 mM		0.5551 mL	2.7753 mL	5.5506 mL
	10 mM		0.2775 mL	1.3877 mL	2.7753 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

1. Add each solvent one by one: PBS
 Solubility: 25 mg/mL (69.38 mM); Clear solution; Need ultrasonic and warming

BIOLOGICAL ACTIVITY

Description

MS023 dihydrochloride is a potent, selective, and cell-active inhibitor of human type I protein arginine methyltransferases (PRMTs) inhibitor, with IC₅₀s of 30, 119, 83, 4 and 5 nM for PRMT1, PRMT3, PRMT4, PRMT6, and PRMT8, respectively^[1].

IC₅₀ & Target

PRMT1	PRMT3	PRMT6	PRMT8
-------	-------	-------	-------

In Vitro

MS023 (1-1000 nM; 48 hours) inhibits PRMT1 methyltransferase activity in MCF7 cells^[1].
 ?MS023(1-1000 nM; 20 hours) inhibits PRMT6 methyltransferase activity in HEK293 cells^[1].
 MCE has not independently confirmed the accuracy of these methods. They are for reference only.
 Western Blot Analysis^[1]

Cell Line:	MCF7 and HEK293 cells
------------	-----------------------

	Concentration:	1.4, 4, 12, 37, 111, 333, and 1000 nM
	Incubation Time:	48 hours for MCF7 cells; 20 hours for HEK293 cells
	Result:	Treatment potently and concentration-dependently reduced cellular levels of H4R3me2a (IC ₅₀ =9±0.2 nM). Treatment concentration-dependently reduced the H3R2me2a mark (IC ₅₀ =56±7 nM).
In Vivo	Administration of MS023 (160 mg/kg, i.p) in combination with PKC412 (100 mg/kg, i.g.) blocks MLL-r acute lymphoblastic leukemia (ALL) propagation by inhibiting maintenance of functional MLL-r ALL-initiating cells ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	
	Animal Model:	NOD-scid IL2Rgnull (NSG) mice bearing primary MLL-r ALL cells ^[2]
	Dosage:	160 mg/kg
	Administration:	Intraperitoneal injection; PKC412 (100 mg/kg, i.g.), MS023 (160 mg/kg, i.p), or a combination for 4 weeks
	Result:	Combinatorial treatment extended survival of leukemic mice relative to single treatments.

CUSTOMER VALIDATION

- Cell Rep. 2021 Sep 21;36(12):109731.
- Acta Pharm Sin B. 22 October 2021.
- Oncogenesis. 2022 Aug 8;11(1):45.
- Acta Pharmacol Sin. 2021 Apr 13.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Eram MS, et al. A Potent, Selective, and Cell-Active Inhibitor of Human Type I Protein Arginine Methyltransferases. ACS Chem Biol. 2016 Mar 18;11(3):772-81.
- [2]. Yinghui Zhu, et al. Targeting PRMT1-mediated FLT3 methylation disrupts maintenance of MLL-rearranged acute lymphoblastic leukemia. Blood. 2019 Oct 10;134(15):1257-1268.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA