



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Ilginatinib hydrochloride

Cat. No.:	HY-19631B
CAS No.:	1239358-85-0
Molecular Formula:	C ₂₁ H ₂₁ ClFN ₇
Molecular Weight:	425.89
Target:	JAK
Pathway:	Epigenetics; JAK/STAT Signaling; Stem Cell/Wnt
Storage:	4°C, sealed storage, away from moisture
	* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro	DMSO : ≥ 35 mg/mL (82.18 mM) H ₂ O : 2 mg/mL (4.70 mM; Need ultrasonic) * "≥" means soluble, but saturation unknown.				
	Preparing Stock Solutions	<div>Solvent Mass Concentration</div>	1 mg	5 mg	10 mg
		1 mM	2.3480 mL	11.7401 mL	23.4802 mL
		5 mM	0.4696 mL	2.3480 mL	4.6960 mL
		10 mM	0.2348 mL	1.1740 mL	2.3480 mL
		Please refer to the solubility information to select the appropriate solvent.			
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (5.87 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	Ilginatinib hydrochloride (NS-018 hydrochloride) is a highly active and orally bioavailable JAK2 inhibitor, with an IC ₅₀ of 0.72 nM, 46-, 54-, and 31-fold selectivity for JAK2 over JAK1 (IC ₅₀ , 33 nM), JAK3 (IC ₅₀ , 39 nM), and Tyk2 (IC ₅₀ , 22 nM).			
IC ₅₀ & Target	JAK2 0.72 nM (IC ₅₀)	Tyk2 22 nM (IC ₅₀)	JAK1 33 nM (IC ₅₀)	JAK3 39 nM (IC ₅₀)
In Vitro	Ilginatinib hydrochloride (NS-018 hydrochloride) is a highly active JAK2 inhibitor, with an IC ₅₀ of 0.72 nM, 46-, 54-, and 31-fold selectivity for JAK2 over JAK1 (IC ₅₀ , 33 nM), JAK3 (IC ₅₀ , 39 nM), and Tyk2 (IC ₅₀ , 22 nM). Ilginatinib hydrochloride also inhibits Src-family kinases, especially SRC and FYN, and weakly inhibits ABL and FLT3 with 45- and 90-fold selectivity for JAK2, respectively. Ilginatinib hydrochloride shows potent inhibitory activity against cell lines JAK2V617F or MPLW515L mutations or the TEL-JAK2 fusion gene (expressing a constitutively activated JAK2) with IC ₅₀ of 11-120 nM, but has only minimal cytotoxicity against most other hematopoietic cell lines that have no constitutively activated JAK2 ^[1] .			

	<p>Ilginatinib hydrochloride (0.5 μM) preferentially suppresses colony-forming unitgranulocyte/macrophage (CFU-GM) formation from myelodysplastic syndrome (MDS)-derived bone marrow mononuclear cells (BMMNCs). Ilginatinib hydrochloride (1 μM) suppresses the phosphorylation of STAT3 (the downstream kinase of JAK2) in CFU-GM-forming cells from MDS patients^[2].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
In Vivo	<p>Ilginatinib hydrochloride (NS-018 hydrochloride) (12.5, 25, 50, 100 mg/kg, p.o.) potently prolongs the survival of mice and reduces splenomegaly in a mouse Ba/F3-JAK2V617F disease model^[1].</p> <p>Ilginatinib hydrochloride (25, 50 mg/kg, p.o.) significantly reduces leukocytosis, hepatosplenomegaly and extramedullary hematopoiesis, improves nutritional status, and prolongs survival in JAK2V617F transgenic mice^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

PROTOCOL

Cell Assay ^[2]	<p>Bone marrow mononuclear cells (BMMNCs) from healthy volunteers and myelodysplastic syndrome (MDS) patients are incubated in MethoCult GF H4434 methylcellulose medium containing various hematopoietic cytokines at 1.0×10^5 cells/mL with or without Ilginatinib (NS-018) at 37°C in a humidified atmosphere of 5% CO₂. Commercially available purified normal human CD34-positive (CD34⁺) BM cells are used as a control. Burst-forming unit-erythroid (BFU-E) and colonyforming unit-granulocyte/macrophage (CFU-GM) colonies are counted under an inverted microscope on day 14 of culture^[2].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
Animal Administration ^[1]	<p>Mice^[1]</p> <p>Female BALB/c nude mice are placed in blanket cages in an environment maintained at 21-25°C and 45-65% relative humidity, with artificial illumination for 12 h and a ventilation frequency of at least 15 times/h. They are allowed free access to food pellets and tap water. Ba/F3-JAK2V617F cells (10^6 per mouse) are inoculated intravenously into 7-week-old mice. Administration of vehicle (0.5% methylcellulose) or Ilginatinib (NS-018) twice daily by oral gavage begins the day after cell inoculation. Survival is monitored daily, and moribund mice are humanely killed and their time of death is recorded for purposes of survival analysis. In a parallel study, all mice are humanely killed after 8 days of administration, and their spleens are removed and weighed^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

REFERENCES

- [1]. Nakaya Y, et al. Efficacy of NS-018, a potent and selective JAK2/Src inhibitor, in primary cells and mouse models of myeloproliferative neoplasms. *Blood Cancer J.* 2011 Jul;1(7):e29.
- [2]. Kuroda J, et al. NS-018, a selective JAK2 inhibitor, preferentially inhibits CFU-GM colony formation by bone marrow mononuclear cells from high-risk myelodysplastic syndrome patients. *Leuk Res.* 2014 May;38(5):619-24.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA