



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

STING agonist-1

Cat. No.:	HY-19711
CAS No.:	702662-50-8
Molecular Formula:	C ₂₁ H ₁₆ ClFN ₂ O ₃ S
Molecular Weight:	430.88
Target:	STING; Virus Protease
Pathway:	Immunology/Inflammation; Anti-infection
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 75 mg/mL (174.06 mM; Need ultrasonic)				
		Mass			
		Solvent	1 mg	5 mg	10 mg
		Concentration			
	Preparing Stock Solutions	1 mM	2.3208 mL	11.6042 mL	23.2083 mL
		5 mM	0.4642 mL	2.3208 mL	4.6417 mL
		10 mM	0.2321 mL	1.1604 mL	2.3208 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (5.80 mM); Clear solution 2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (5.80 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	STING agonist-1 (G10) is human-specific STING agonist that elicits antiviral activity against emerging Alphaviruses. G10 potently blocks replication of Alphavirus species Venezuelan Equine Encephalitis Virus (VEEV) with IC ₉₀ of 24.57 μM ^[1] .
IC ₅₀ & Target	IC ₉₀ : 24.57 μM (VEEV) ^[1]
In Vitro	G10 induces IFN/IRF3- but not NF-κB-dependent transcription in human fibroblasts ^[1] . G10 is an indirect activator of human STING-dependent phenotypes ^[1] . G10-mediated IRF3 activation and anti-Alphaviral activity occur independently of IPS-1/MAVS-dependent signaling ^[1] . STING is required for G10-mediated IRF3 activation, gene expression, and anti-Alphaviral activity ^[1] . G10 induces STING-dependent synthesis and secretion of bioactive interferon ^[1] .

G10 induces innate antiviral mRNA expression in primary human cells^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cancer Immunol Res. 2023 May 3;11(5):583-599.
- Patent. US20200268864A1.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Sali TM, et al. Characterization of a Novel Human-Specific STING Agonist that Elicits Antiviral Activity Against Emerging Alphaviruses. PLoS Pathog. 2015 Dec 8;11(12):e1005324.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA