



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

LJI308

Cat. No.:	HY-19713
CAS No.:	1627709-94-7
Molecular Formula:	C ₂₁ H ₁₈ F ₂ N ₂ O ₂
Molecular Weight:	368.38
Target:	Ribosomal S6 Kinase (RSK); YB-1
Pathway:	MAPK/ERK Pathway; Cell Cycle/DNA Damage
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 25 mg/mL (67.86 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM	2.7146 mL	13.5729 mL	27.1459 mL	
		5 mM	0.5429 mL	2.7146 mL	5.4292 mL	
		10 mM	0.2715 mL	1.3573 mL	2.7146 mL	
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% corn oil					
	Solubility: ≥ 2.5 mg/mL (6.79 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	LJI308 is a potent pan-ribosomal S6 kinase (RSK) inhibitor, with IC ₅₀ s of 6 nM, 4 nM, and 13 nM for RSK1, RSK2, and RSK3, respectively. LJI308 inhibits the phosphorylation of RSK (T359/S363) and YB-1 (S102) after irradiation, treatment with EGF, and in cells expressing a KRAS mutation ^{[1][2]} .		
IC ₅₀ & Target	RSK1	RSK2	RSK3
In Vitro	LJI308 inhibits S6K1 with an IC₅₀ of 0.8 μM^[1]. LJI308 inhibits YB-1 phosphorylation in CRC cells at concentrations of 5 to 25 μM. In a dose kinetics experiment, LJI308, starting at 2.5 μM, inhibits YB-1 phosphorylation in the KRAS mutated TNBC cell line MDA-MB-231 by approximately 86%. LJI308 effectively blocks RSK and YB-1 phosphorylation after EGF stimulation and after irradiation in KRAS wild-type HBL-100 cells^[2]. LJI308 (1-10 μM; 96 hours) decreases cell viability by up to 90%^[3].		

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Viability Assay^[3]

Cell Line:	HTRY-LT cell lines
Concentration:	1-10 μ M
Incubation Time:	96 hours
Result:	Decreased cell viability by up to 90%.

CUSTOMER VALIDATION

- Circulation. 2022 Nov 30.
- Exp Hematol Oncol. 2023 Nov 30;12(1):100.
- J Invest Dermatol. 2020 Sep 9;S0022-202X(20)32055-8.
- Oncotarget. 2017 Jun 6;8(23):37633-37645.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Aronchik I, et al. Novel potent and selective inhibitors of p90 ribosomal S6 kinase reveal the heterogeneity of RSK function in MAPK-driven cancers. Mol Cancer Res. 2014 May;12(5):803-12.
- [2]. Lettau K, et al. Simultaneous Targeting of RSK and AKT Efficiently Inhibits YB-1-Mediated Repair of Ionizing Radiation-Induced DNA Double-Strand Breaks in Breast Cancer Cells. Int J Radiat Oncol Biol Phys. 2021;109(2):567-580.
- [3]. Jain R, et al. Discovery of Potent and Selective RSK Inhibitors as Biological Probes. J Med Chem. 2015 Sep 10;58(17):6766-83.
- [4]. Davies AH, et al. Inhibition of RSK with the novel small-molecule inhibitor LJI308 overcomes chemoresistance by eliminating cancer stem cells. Oncotarget. 2015;6(24):20570-20577.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA