



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

BCX 1470 methanesulfonate

Cat. No.:	HY-50875
CAS No.:	217099-44-0
Molecular Formula:	C ₁₅ H ₁₄ N ₂ O ₅ S ₃
Molecular Weight:	398.48
Target:	Complement System
Pathway:	Immunology/Inflammation
Storage:	4°C, sealed storage, away from moisture * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 33.33 mg/mL (83.64 mM)

* "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
	1 mM	2.5095 mL	12.5477 mL	25.0954 mL	
	5 mM	0.5019 mL	2.5095 mL	5.0191 mL	
	10 mM	0.2510 mL	1.2548 mL	2.5095 mL	

Please refer to the solubility information to select the appropriate solvent.

In Vivo

1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline

Solubility: ≥ 2.5 mg/mL (6.27 mM); Clear solution

2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)

Solubility: ≥ 2.5 mg/mL (6.27 mM); Clear solution

3. Add each solvent one by one: 10% DMSO >> 90% corn oil

Solubility: ≥ 2.5 mg/mL (6.27 mM); Clear solution

BIOLOGICAL ACTIVITY

Description	BCX 1470 methanesulfonate inhibits the esterolytic activity of factor D (IC ₅₀ =96 nM) and C1s (IC ₅₀ =1.6 nM), 3.4- and 200-fold better, respectively, than that of trypsin.
IC ₅₀ & Target	IC ₅₀ : 96 nM (Factor D); 1.6 nM (C1s); 326 nM (Trypsin)
In Vitro	BCX 1470 methanesulfonate is serine protease inhibitor. BCX 1470 methanesulfonate blocks the esterolytic and hemolytic activities of the complement enzymes C1s and factor D in vitro, also blocked development of RPA-induced edema in the rat. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Szalai AJ, et al. The Arthus reaction in rodents: species-specific requirement of complement. J Immunol. 2000 Jan 1;164(1):463-8.
- [2]. Szalai AJ, et al. The Arthus reaction in rodents: species-specific requirement of complement. J Immunol. 2000 Jan 1;164(1):463-8.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA