



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

N-desmethyl Enzalutamide

Cat. No.:	HY-70002A
CAS No.:	1242137-16-1
Molecular Formula:	C ₂₀ H ₁₄ F ₄ N ₄ O ₂ S
Molecular Weight:	450.41
Target:	Androgen Receptor
Pathway:	Vitamin D Related/Nuclear Receptor
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (222.02 mM; Need ultrasonic)				
	Preparing Stock Solutions	Solvent Concentration Mass	1 mg	5 mg	10 mg
		1 mM	2.2202 mL	11.1010 mL	22.2020 mL
		5 mM	0.4440 mL	2.2202 mL	4.4404 mL
		10 mM	0.2220 mL	1.1101 mL	2.2202 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (5.55 mM); Clear solution				
	2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (5.55 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	N-desmethyl Enzalutamide is the active metabolite of Enzalutamide. N-desmethyl Enzalutamide is the active metabolite of Enzalutamide. N-desmethyl Enzalutamide demonstrates primary and secondary pharmacodynamics of similar potency to Enzalutamide and circulates at approximately the same plasma concentrations as enzalutamide ^[1] .
IC ₅₀ & Target	Androgen-receptor ^[1]
In Vivo	N-desmethyl Enzalutamide is an active metabolite that is thought to contribute to the clinical effects of Enzalutamide because it demonstrates primary and secondary pharmacodynamics of similar potency to Enzalutamide and circulates at approximately the same plasma concentrations as Enzalutamide. The carboxylic acid metabolite is pharmacologically inactive and circulates at approximately 25 % lower plasma concentrations than Enzalutamide ^[1] .

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Gibbons JA, et al. Pharmacokinetic Drug Interaction Studies with Enzalutamide. Clin Pharmacokinet. 2015 Oct;54(10):1057-69.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA