



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Enzalutamide carboxylic acid

Cat. No.:	HY-70002B
CAS No.:	1242137-15-0
Molecular Formula:	C ₂₀ H ₁₃ F ₄ N ₃ O ₃ S
Molecular Weight:	451.39
Target:	Drug Metabolite
Pathway:	Metabolic Enzyme/Protease
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 125 mg/mL (276.92 mM; Need ultrasonic)				
	Preparing Stock Solutions	Solvent Concentration Mass	1 mg	5 mg	10 mg
		1 mM	2.2154 mL	11.0769 mL	22.1538 mL
		5 mM	0.4431 mL	2.2154 mL	4.4308 mL
		10 mM	0.2215 mL	1.1077 mL	2.2154 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (4.61 mM); Clear solution				
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (4.61 mM); Clear solution				
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (4.61 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	Enzalutamide carboxylic acid (MDV3100 carboxylic acid) is an inactive metabolite of Enzalutamide (MDV3100). Enzalutamide is an androgen receptor (AR) antagonist ^[1] .
In Vitro	Enzalutamide is converted into its major metabolites, N-desmethyl enzalutamide and carboxylic acid enzalutamide, by cytochrome P450 (CYP) 3A4/5 and CYP2C8, respectively. N-desmethyl enzalutamide has clinically relevant anti-androgen capacities similar to enzalutamide, whereas carboxylic acid enzalutamide is inactive ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Clin Transl Med. 2022 Apr;12(4):e797.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. van Nuland M, et al. Exposure-Response Assessment of Enzalutamide and Its Major Metabolites in a Real-World Cohort of Patients with Metastatic Castration-Resistant Prostate Cancer. Pharmacotherapy. 2019 Dec;39(12):1137-1145.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA