



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Quetiapine-d₄-1 fumarate

Cat. No.:	HY-B0031S5
CAS No.:	2820085-77-4
Molecular Formula:	C ₂₁ H ₂₁ D ₄ N ₃ O ₂ S.1/2C ₄ H ₄ O ₄
Molecular Weight:	445.57
Target:	Dopamine Receptor; 5-HT Receptor; Isotope-Labeled Compounds
Pathway:	GPCR/G Protein; Neuronal Signaling; Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	<p>Quetiapine-d₄-1 fumarate is deuterated labeled Quetiapine (hemifumarate) (HY-B0031). Quetiapine hemifumarate is a 5-HT receptors agonist with a pEC₅₀ of 4.77 for human 5-HT_{1A} receptor. Quetiapine hemifumarate is a dopamine receptor antagonist with a pIC₅₀ of 6.33 for human D₂ receptor. Quetiapine hemifumarate has moderate to high affinity for the human D₂, HT_{1A}, 5-HT_{2A}, 5-HT_{2C} receptor with pK_s of 7.25, 5.74, 7.54, 5.55. Antidepressant and anxiolytic effects^[1].</p>
In Vitro	<p>Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs^[1].</p> <p>Quetiapine (<100?μM; 24?hours) has no significant effect on cell viabilities^[3].</p> <p>Quetiapine (10?μM) inhibits NO release, which increased by LPS (0.1-100 ng/mL) in concentration-dependent manner^[3].</p> <p>Quetiapine (10?μM) also inhibits TNF-α synthesis^[3].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
In Vivo	<p>Quetiapine (10?mg/kg/day; ingested) can alleviate the recruitment and activation of microglia and promote myelin repair in Cuprizone (CPZ)-induced chronic mouse model of demyelination^[2].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

REFERENCES

- [1]. Cross AJ, et al. Quetiapine and its metabolite norquetiapine: translation from in vitro pharmacology to in vivo efficacy in rodent models. Br J Pharmacol. 2016 Jan;173(1):155-66.
- [2]. Hanzhi Wang, et al. Quetiapine Inhibits Microglial Activation by Neutralizing Abnormal STIM1-Mediated Intercellular Calcium Homeostasis and Promotes Myelin Repair in a Cuprizone-Induced Mouse Model of Demyelination. Front Cell Neurosci. 2015 Dec 21;9:492.
- [3]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA