



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Pizotyline-d₃

Cat. No.:	HY-B0115S
Molecular Formula:	C ₁₉ H ₁₈ D ₃ NS
Molecular Weight:	298.46
Target:	5-HT Receptor; Isotope-Labeled Compounds
Pathway:	GPCR/G Protein; Neuronal Signaling; Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Pizotyline-d ₃ is deuterated labeled Pizotifen (HY-B0115). Pizotifen (Pizotyline) is a potent 5-HT ₂ receptor antagonist, with a high affinity for 5-HT _{1C} binding site.
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs^[1]. Pizotifen (BC-105) is a potent 5-HT₂ receptor antagonist, with a high affinity for 5-HT_{1C} binding site^[2]. Pizotifen is an antidepressant 5-HT_{2A} receptor antagonist and has the capacity to inhibit serotonin-enhanced ADP-induced platelet aggregation^[3]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	The weights of the fetuses are significantly reduced by all administered doses of Pipethiadene and Pizotifen (BC-105); the weights of the placentas are significantly reduced after 0.6 and 1.2 mg/kg Pipethiadene and only after the middle dose of Pizotifen. The means of the implantations, live, dead fetuses, resorptions and the occurrence of external, skeletal and visceral anomalies do not differ from the control group. The number of chromosome aberrations in the bone marrow cells of treated mice does not differ significantly from the negative control group. The micronucleus test reveals no elevation in the frequency of micronuclei as compared to the control group. After the two higher doses of both Pipethiadene and Pizotifen (BC-105) maleate, the mitotic indices are lower than in the control group^[4]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Mylecharane EJ, et al. 5-HT₂ receptor antagonists and migraine therapy. J Neurol. 1991;238 Suppl 1:S45-52.
- [2]. Lin OA, et al. The antidepressant 5-HT_{2A} receptor antagonists pizotifen and cyproheptadine inhibit serotonin-enhanced platelet function. PLoS One. 2014 Jan 23;9(1):e87026.
- [3]. Ujházy E, et al. Teratological and cytogenetical evaluation of two antihistamines (pipethiadene and pizotifen maleate) in mice. Agents Actions. 1988 Apr;23(3-4):376-8.
- [4]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. Ann Pharmacother. 2019 Feb;53(2):211-216.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA